

Antibiotic Susceptibility Profile of Bacteria Isolated from Wound Sepsis Patients from Immanuel Hospital Eket, Akwa Ibom State, Nigeria

¹Christopher, M. A., ²Nyoyoko, V. F., ¹Ibanga, I. A., ¹Owowo, E. E. and ¹Peter, P. U.

¹Department of Microbiology, Akwa Ibom State University, Mkpato-Enin, Nigeria

²Department of Microbiology, University of Nigeria, Nsukka, Nigeria

Corresponding author: meritony27@gmail.com:08034857589

Abstract: Antibiotic Susceptibility Profile of Bacteria Isolated from Wound Sepsis Patients was study. The study aimed at assessing the antibiotic susceptibility of some bacteria isolated from wound sepsis of patients attending Immanuel General Hospital Eket. Random sampling technique was employed in this study. One hundred (100) wound swabs were obtained and cultured on Blood agar and MacConkey agar. Most Bacterial isolates from the wound were susceptible to most of the antibiotics with few of them being resistant. The prevalence of bacteria isolated from wound swabs were *Staphylococcus aureus* (42.8%), *Escherichia coli* (14.3%), *Pseudomonas aeruginosa* (14.3%), *Streptococcus pyogenes* (11.4%), *Klebsiella pneumoniae*, (8.6%) and *Proteus* sp. (8.6%). The percentage susceptibility results showed that *Staphylococcus aureus* was highly susceptible to Rifampicin (96%), Streptomycin (92%) and Levofloxacin (88%). *Streptococcus pyogenes*, exhibited 100% susceptibility to Ciprofloxacin, Chloramphenicol and Levofloxacin but strongly resisted Amoxicillin (75%). *Escherichia coli* showed high susceptibility to Septrin (90%), Augmentin (90%), Tarivid (80%) and Nalidixic acid (70%). *Pseudomonas aeruginosa* and *Proteus* spp. isolates were resistant to Ciprofloxacin, Gentamycin. *Klebsiella pneumoniae* showed resistance to Ciprofloxacin (75%), Augmentin (75%) and Streptomycin (62%). The demographic factor; sex, age, occupation, area of domicile of patients was associated with the occurrence of bacteria from the wound sepsis. The frequency of single and multiple drug resistance is highly alarming and is a growing threat to the control of infectious diseases globally.

Keywords: Antibiotic, Susceptibility, Bacteria, Wound Sepsis

INTRODUCTION

A wound is a type of injury which happens relatively quickly in which skin is torn, cut or punctured (an open wound or where blunt force trauma causes a contusion (a closed wound) (Adegoke *et al.*, 2010; Siddiqui and Bernstein 2010; Ohalet *et al.*, 2012). The risk of wound infection increases with the degree of contamination and it has been estimated that about 50% of wounds contaminated with bacteria become clinically infected (Omole and Stephen 2014), (Sahu *et al.*, 2011). Contamination of wound occurs when non-replicating bacteria enters the wound (Sahu *et al.*, 2011; Tom *et al.*, 2018). Contamination can also occur when the bacteria begin replicating and adhere to the wound site but do not cause tissue damage (Goldstein *et al.*, 1996). It can also occur when the number of bacterial is greatly increased and begins to overwhelm the host immune system (Fonder *et al.*, 2008). During this stage, the granulation bed in the wound appears unhealthy example

atrophied, deep red or grey discoloration, with increased discharged but there is no sign of invasion of the surrounding tissue (Sani *et al.*, 2012; Fonder *et al.*, 2008; Janet *et al.*, 2005). Wound infections are classified on a continuum; contaminated, colonized, local infection, spreading infection, and systemic infection (sepsis). Infections of the skin and soft tissue either due to trauma, surgery, or burns may result in the generation of exudates composed of dead leucocytes, cellular debris, and necrotic tissues (Dryden, 2010). Chronic wounds can be colonized on the surface by a wide range of organisms (Matsuura and Barg, 2013). Wound infection is characterized by the presence of pus in lesions with pyrexia, pain and in duration. Infection occurs when virulence factors produced by the microorganisms overwhelms the host natural resistance (Bowler *et al.*, 2010). The contamination or mere presence of pathogenic organisms in wound without local or systemic tissue involvement may not result in infection (Sule *et al.*, 2002).

The wound environment is conducive for bacterial growth. The organisms can cause destruction if left untreated (Austrian Wound Management Association, 2011). Bacterial wound infections are important because they can slow down the healing process, lead to wound breakdown, prolonged hospital stay and increase in the cost of treatment (Pondei *et al.*, 2013; Duce *et al.*, 2010). Common bacterial pathogens associated with wound infection include *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Streptococcus pyogenes*, *Proteus spp.*, *Bacteriodes spp.*, *Clostridium*, and *Enterococcus spp.* (Dani *et al.*, 2012). These organisms exhibit natural resistance to many antibiotics and antiseptics in which they may survive for long periods, and may even multiply in the presence of minimal nutrients and have the ability to colonize traumatised skin (Richard *et al.*, 1994; Olayinka *et al.*, 2004). Wounds are the third most frequent nosocomial infections (Duce *et al.*, 2010). In developing and resource-poor countries, traumatic and surgical site infections are reasons for high morbidity and mortality rates (Egbe *et al.*, 2011). The likelihood of a wound becoming infected is related to the number and virulence of the infecting microorganism and the ability of the host to resist infection (ICU, 2016) of these opportunistic pathogens form part of the host's normal microbiota, or are found in damp environmental sites or on hospital equipments and medicament (Olayinka *et al.*, 2004). Once they gain access into the body, they develop mechanisms to exploit the host for continuous survival and dissemination (Tom *et al.*, 2018). Patients with wound complication arising from the dissemination of pathogenic microorganisms tend to be associated with bacteraemia, septicaemia, shock and prolonged hospital stay with an increasing chance of developing drug resistant infections. Drug resistance leads to prolonged epidemics (Nita *et al.*, 2018), and consequently, an unattended wound-site, being the most vulnerable point

of entry of pathogenic bacteria could be difficult to treat if a multi-drug resistant strain is implicated. This development is worrisome with a resultant increase in morbidity, mortality and cost not only to patients and their relatives but including hospital management (Baba *et al.*, 2016).

MATERIALS AND METHODS

Study Area

The samples were processed at Akwa Ibom State University microbiology laboratory, Ikot Akpaden, Mkpato Enin Local Government Area, as they were obtained from patients attending Immanuel General Hospital Eket. Eket local Government occupies the South-Central portion of Akwa Ibom State territorial expanses spanning Northwards between latitude 4°33' and 4°45' and Eastward between Longitudes 7°52' and 5°02'. It is bounded on the north by Nsit Ubium Local Government Area, on the east by Esit Eket local Government Area, on the West by Onna Local Government Area and on the South by Ibeno Local Government Area / Bight of Bonny. Immanuel Hospital named after a Lutheran Church is a missionary hospital which during the Civil war gave room to government involvement which provided enough medical facilities and equipments able to treat patients.

Sample Collection

Random sampling method was used to select the study participants which comprised of 100 patients with various types of wound infections. Data for the study were generated by administering structured questionnaire and by collecting wound swabs. The structured Questionnaires were administered to generate information on the age, gender, occupation, educational qualification and area of domicile (resident). Swab sticks were used to collect different types of wound samples (Trauma, Postoperative wound, Abscess, Ulcer, Burn wounds and Diabetic foot ulcer and location Leg, Abdomen, Hand, Foot, Ankle, Back, Head and neck) from patients with infected wounds.

Isolation of Bacteria from Wound Swabs

A total of 100 wound swab samples were collected. The swab specimens were streaked on blood agar and MacConkey agar and then placed in the incubator. The swab specimens were incubated for 24 hours at 37°C; the plates were examined for growth and different pattern of hemolysis. The plates were then sub-cultured into MacConkey agar, and pure cultures were obtained and transferred into Nutrient agar slants for further identifications (Cheesbrough, 2010).

Identification of Bacterial Isolates

Identifications were carried out by sub-culturing from the Nutrient agar slants into freshly prepared MacConkey and Blood agar in sterile petri dishes using the streaking method. The pure culture of the isolates were subjected to biochemical tests, namely: Catalase, Coagulase, Indole, Citrate utilization, Urease, Methyl red-Voges Proskauer, Motility and Sugar fermentation for identification after Gram staining which was used to test for the organisms purity (Cheesbrough, 2010).

Antibiotics Susceptibility Test

The antibacterial susceptibility testing was done using Kirby-Bauer NICCS modified disc agar diffusion technique: all procedure was done under aseptic technique. Isolated bacteria were subculture in normal saline for 3 hours interval to obtain a solution with turbidity equal to 0.5 McFarland standards. Sterile swab stick was used to evenly spread the organism across the Muller Hinton agar plate and allowed to dry. The impregnated disks were carefully picked with sterile forceps and carefully placed on the inoculated Muller Hinton agar plate. The plates were incubated at 37°C for 24 hours. The diameters of the zones of inhibition were measured and recorded for the urine samples that showed zones of inhibition. The zones of inhibition were measured using a meter rule. All measurements were recorded in millimeters. The Sensitivity pattern of the isolates to Tarivid, Reflacine, Ciprofloxacin, Augmentin, Gentamycin, Streptomycin, Ceporex, Nalidixic acid,

Septtrin, Norfloxacin, Amoxil, Rifampin Erthromycin, Chloramphenicol, Ampiclox, were determined. Isolates were divided into three groups based on the zone of inhibition produced by the antibiotic disc; susceptible, intermediately susceptible and resistant according to the Clinical and Laboratory Standards Institute (CLSI) guideline; Performance Standards for Antimicrobial Susceptibility Testing (2007).

Statistical Analysis

This was analyzed using Chi- Square test of association between the socio - demographic factors and the occurrence of bacteria in the wound sepsis of the patients.

RESULTS

Table 1 showed the percentage occurrence of the different bacteria isolated where *Staphylococcus aureus* was the most predominant with 15 (42.8%) isolates. This is followed by *Escherichia coli* and *Pseudomonas aeruginosa* with 5 (14.3%) isolates each, *Streptococcus pyogenes* with 4 (11.4%) and the least were *Klebsiella pneumoniae* and *Proteus spp* with 3(8.6%) isolates each.

Table 2 shows Susceptibility of Gram negative isolates to some antibiotics *Escherichia coli* had high susceptibility to Septtrin (90%), Augmentin (90%), Tarivid (80%) and Nalidixic acid (70%). *Klebsiella pneumoniae* showed susceptibility to most of the antibiotics, but resisted Ciprofloxacin, Augmentin and Streptomycin. *Pseudomonas aeruginosa* and *Proteus spp* isolates showed susceptibility to all the antibiotics, but resisted Ciprofloxacin, Gentamycin and Septtrin (Table 2).

Table 3 showed the percentage susceptibility of Gram positive isolates to some antibiotics were *Staphylococcus aureus* isolates were highly susceptible to Rifampicin (96%), Streptomycin (92%), and Levofloxacin (88%). In the case of *Streptococcus pyogenes*, the isolates exhibited 100% susceptibility to Ciprofloxacin, Chloramphenicol, and Levofloxacin and strongly resisted Amoxil (75%). Table 4

showed the Biochemical Characteristics of Bacteria Isolated from Wound Sepsis in Patient. The demographic factor; sex, age, occupation, area of domicile, site of infection and types of wound of patients are presented in Table 5.

Table 1: Percentage Occurrence of the Bacterial Isolates from Wound Sepsis of Patients attending Immanuel General Hospital Eket

Bacteria	Total number	Percentage (%)
<i>Staphylococcus aureus</i>	15	42.8
<i>Escherichia coli</i>	5	14.3
<i>Pseudomonasaeruginosa</i>	5	14.3
<i>Streptococcus pyogenes</i>	4	11.4
<i>Klebsiellapneumonia</i>	3	8.6
<i>Proteusspp</i>	3	8.6
Total	35	100.0

Table 2: Percentage Susceptibility of Gram-Negative Isolates to some Antibiotics

ANTIBIOTICS	<i>Escherichia coli</i> (n=5)%	<i>Klebsiella pneumoniae</i> (n=3)%	<i>Pseudomonas aeruginosa</i> (n=5)%	<i>Proteus spp</i> (n=3)%
Tarivid	80	88	66	66
Reflacine	70	75	100	100
Ciprofloxacin	40	25	0	0
Augmentin	90	25	66	100
Gentamycin	30	50	0	33
Streptomycin	60	38	100	66
Ceporex	50	88	66	100
Nalidixic acid	70	75	100	66
Seprtin	90	50	33	33
Amplicin	50	75	66	100

Table 3: Percentage Susceptibility of Gram-Positive Isolates to some Antibiotics

Antibiotics	<i>Staphylococcus aureus</i> (n=15)%	<i>Streptococcus pyogenes</i> (n=4)%
Ciprofloxacin	65	100
Norfloxacin	69	50
Gentamycin	19	75
Amoxil	38	25
Streptomycin	92	50
Rifampicin	96	75
Erythromycin	18	75
Chloramphenicol	57	100
Ampiclox	23	75
Levofloxacin	88	100

Table 4: Biochemical Characteristics of Bacteria Isolated from Wound Sepsis in Patient Attending Immanuel General Hospital Eket, Akwa Ibom State, Nigeria

Isolates	Catalase	Coagulase	Indole	Methyl	Voges proskauer	Urease	Motility	Sugar fermentation				Gram staining	Most probable micro- organism
								Glucose	Sucrose	lactose	Gas		
1	+	+	-	+	+	+	-	+	+	+	-	Positive cocci in chains	<i>Staphylococcus aureus</i>
2	+	-	+	+	-	-	+	+	Variable	+	+	Negative rods	<i>Escherichia coli</i>
3	+	-	-	-	-	-	+	+	-	-	-	Negative rods	<i>Pseudomonas aeruginosa</i>
4	-	-	-	-	-	-	-	+	+	+	-	Positive cocci	<i>Streptococcus pyogenes</i>
5	+	-	-	-	+	+	-	+	+	+	+	Negative rods	<i>Klebsiella pneumoniae</i>
6	+	-	-	+	-	+	+	+	-	-	+	Negative rods	<i>Proteus spp</i>

Keywords: + (positive), - (negative)

T

TABLE 5: The Demographic Factor of Patients with the Wound Sepsis

Variable	No Studied	No Infected	% Infected	X ²	P-value		
Age							
1-10	7	7	100	16.4751	0.033		
11-20	19	19	100				
21-30	24	24	100				
31-40	17	17	100				
41-50	15	15	100				
51-60	10	10	100				
61-70	8	8	100				
Sex							
Male	65	65	100	9.0000	0.083		
Female	35	35	100				
Occupation							
Farmer	30	30	100	16.1566	0.075		
Cyclist	18	18	100				
Business owner	8	8	100				
Non-employed student	16	16	100				
Student	14	14	100				
Employed	14	14	100				
Area of Domicile							
Urban	68	68	100			12.9600	0.083
Rural	32	32	100				
Site of Infection							
Foot	28	28	100	51.7445	0.017		
Leg	25	25	100				
Hand	25	25	100				
Shoulder	8	8	100				
Ankle	5	5	100				
Knee	6	6	100				
Finger	3	3	100				
Types of Wounds							
Diabetic	28	28	100	19.6361	0.205		
Ulcer	20	20	100				
Burn	13	13	100				
Post Operation	20	20	100				
Abscess	15	15	100				
Trauma ulcer	4	4	100				

X²=Chi-square

DISCUSSION

The high prevalence of *Staphylococcus aureus* as revealed by the study could be related to the fact that this organism is an endogenous microbial flora. Infections with these bacteria may also be due to contamination from the environment e.g. contamination of surgical instruments. With the disruption of the natural skin barrier, *Staphylococcus aureus*, which is a common bacterium on surfaces, easily find their way into wounds. Similarly, high percentage of microbial growth was reported in Minna (Baba *et al.*, 2016), Ekpoma (Emele *et al.*, 1999), India (86-100%) and Pakistan (98%) (Basu *et al.*, 2009). Most of the isolates identified in this study had mixed with other bacterial species and some of these have shown to be resistant to many antibiotics, and this indicates the high contamination of wounds in our hospitals. In the present study, the most commonly isolated bacteria from wounded patients were *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa*. The reasons for this high prevalence may be due to factors associated with the acquisition of nosocomial pathogens in patients on long admission, thus complicating illnesses, prior to the administration of antimicrobial agents. The result of percentage prevalence of the bacteria isolates from this study also indicated that the rate of isolation of Gram-positive bacteria was more than that of Gram-negative bacteria. This finding negates the report of Obritsch *et al.* (2004), who reported that the rate of Gram Negative bacteria isolation from wound was more than twice that of Gram Positive, and it was noticed that *Klebsiella spp.* was the pathogen less isolated followed by *Pseudomonas aeruginosa* and *Staphylococcus aureus*.

Pseudomonas aeruginosa and *Proteus spp* were found to be highly resistant to Gentamicin and Ciprofloxacin. High resistance of the isolates to antibiotics may be due to practicing self-medication, lack of diagnostic laboratory services or unavailability of guideline regarding the

selection of drugs thereby leading to inappropriate use of antibiotics. This finding is supported by the investigation of Obritsch *et al.* (2004), who reported that more than 72% of the Gram negative isolates of wound were resistant to Gentamycin, a commonly used antibiotic against Gram positive infections. The study findings indicated that *Staphylococcus aureus* isolates are highly susceptible to Rifampicin (96%), Streptomycin (92%), and Levofloxacin (88%). In the case of *Streptococcus pyogenes*, the isolates exhibited 100% susceptibility to Ciprofloxacin, Chloramphenicol, and Levofloxacin and strongly resisted Amoxil. *E. coli* showed high susceptibility to Seprin,–Augmentin, Tarivid and Nalidixic acid.

Pseudomonas aeruginosa showed susceptibility to most of the antibiotics, but resisted Ciprofloxacin, Augmentin and Streptomycin. This might be as a result of inappropriate use of antimicrobial agents (antibiotics) and the intrinsic ability of the isolates to survive in their environment. This work disagrees with the study done in Ethiopia with average susceptibility of *Staphylococcus aureus* and *Streptococcus spp* isolates (0%) and *E. coli* isolates (4.5%) (Mulu *et al.*, 2012) but is relative to the research carried out in Minna General Hospital Nigeria as reported by Baba *et al.* (2016). The results of antibiotic susceptibility showed that *Streptococcus pyogenes* strongly resisted Amoxil.

The biochemical characterization and differentiation into both Gram Positive and Gram Negative isolate further revealed and confirmed the isolate nutritional and metabolic capabilities so as to know the kind of drug that will be used to targets it as it affects patient outcome. There is an association between the demographic factor; sex, age, occupation, area of domicile of patients and occurrence of bacteria from the wound sepsis.

CONCLUSION

The study identified that most of the bacterial isolates from the wound are

susceptible to most of the antibiotics with few of them being resistant.

This research addresses the microbiological aspect that are critical to the successful management of bacteria in wound and resistant rate of these bacteria to single and multiple antibiotics which is really alarming and a threat to the control of infectious diseases. The demographic factor; sex, age, occupation, area of domicile of patients was

associated with the occurrence of bacteria from the wound sepsis. The frequency of single and multiple drug resistance is highly alarming and is a growing threat to the control of infectious diseases globally.

CONFLICT OF INTEREST

The authors declare no conflict of interest

REFERENCES

- Abe, Y., Kashiwagi, K., Ishida, S., Mineda, K., Yamashita, Y. and Hashimoto, I. (2018). Risk factors for delayed healing at the free anterolateral thigh flap donor site. *Archaic Plastic Surgery*. 45(1):51-57.
- Adegoke, A. A., Tom, M., Okoh, A. I., and Jacob, S. (2010). Studies on multiple antibiotic resistant bacterial isolated from surgical site infection. *Scientific Research and Essay*. 5:3876-3881.
- Ameh, V. O. (2014). Epidemiological studies of canine rabies in Wukari metropolis, Taraba State, Nigeria. *Thesis submitted to the School of Postgraduate Studies, Ahmadu Bello University, Zaria, Nigeria*.
- American Association of Clinical Chemistry (AACC). (2018). *Journal of Applied Laboratory Medicine*, Retrieved May, 2020.
- Amoran, O. E., Sogebi, A. O and Fatugase, O. M. (2013). Rates and risk factors associated with surgical site infections in a tertiary care center in south-western Nigeria. *International Journal of Tropical Disease and Health*, 3(1):25-36.
- Andrew Gonzalez, M. D., Julie Roddick, J. D. and M. P. H. (2018). Open wounds. Open wound, *Health line.com research journal* Medically reviewed in September. CD102766.
- Arabishahi, K. S. and Koohpayezade, J. (2006). Investigation of risk factors for surgical wound infection among teaching hospital in Tehran. *International wound Journal*. 3: 59-62.
- Australian Wound Management Association (AWMA). (2011). Bacterial impact on wound healing: from contamination to infection. *Position Paper*, 1-16.
- Baba, J., Olutimayin A. T., Alalade O. M., Aliyu M. B. and Ndagi G. M. (2016). Isolation and identification of some bacteria in wound sepsis. *Lapal Journal of Applied Sciences*. LAJANS 1(1):104-110.
- Basu, S., Ramchuran, P. T., Bali, S.T., Gulat, I. A. and Shukla, V. (2009). *A Perspective, descriptive study to identify the microbiological profile of chronic wounds in outpatients*. *Ostomy Wound Management*, 55(1):14-20.
- Bree, N. and Stephanie, W. (2020). Cellulitis. Medically reviewed by Debra Sullivan on February 19, 2020.
- Brewer, J. D., Gonzalez, A. B., Baum, C. L., Arpey, C. J., Roenigk, R. K., Otley, C. C. and Erwin, P. J (2016). "Comparison of Sterile vs Nonsterile Gloves in Cutaneous Surgery and Common Outpatient Dental Procedures: A Systematic Review and Meta-analysis". *JAMA Dermatology*. 152 (9):10-14.
- Bowler, P. G., Duerden, B. I. and Armstrong, D. G. (2001). Wound microbiology and associated approaches to wound management. *Clinical microbiological review*, 14 (2): 244-269.
- Campbell, J. L., Coyer, G. M. and Osborne, S. R. (2018). The skin safety model: reconceptualizing skin vulnerability in older patients. *Journal on Nursing Scholarship*. 48(1):14-22.
- Cantlon, C. A., Stemper, M. E., Schwan, W. R., Hoffman, M. A. and Qutaishat, S. S (2006). Significant pathogen isolated

- from surgical site infections at a community hospital in the Midwest. *Journal on Infection Control*, 34: 526-529.
- Cheesebrough, M. (2010). Microbiological District Laboratory Practice in Tropical Countries. Tropical Health Technology, Norfolk. 2nd edition. United Kingdom, Cambridge press. 189.
- Cheryl, C. (2017). Fungal Infections, Moisture-Associated Skin Damage, Skin Care fungi *candida albicans* 3D.LPN, WCC, CWCA, FACCWS, DAPWCA, CLTC.
- Chiller, K., Selkin, B. A. and Murakawa G. J. (2001). Skin microflora and bacterial infections of the skin. *Journal Investigating Dermatology*. 6:170-174.
- Christensen, G. J. and Brüggemann, H. (2014). Bacterial skin commensals and their role as host guardians. *Beneficial Microbes*. 5:201-215.
- CLSI (2007) Performance standards for antimicrobial susceptibility testing; Seventeenth informational supplement. Clinical and Laboratory Standards Institute 27: M100-S17.
- Cone, L. A., Leung, M. M. and Hirschberg, J. (2003). *Actinomyces odontolyticus* bacteremia. *Emergence of Infectious Disease*. 2003 Dec. 9 (12):1629-1632. [Medline].
- Dean, J. (2011). Skin health: prevention and treatment of skin breakdown. *Transverse Myelitis Association Journals*. 11;5.
- Dong, Y. and Speer C. P. (2014). The role of *Staphylococcus epidermidis* in neonatal sepsis: guarding angel or pathogenic devil? *International Journal on Medical Microbiology*. 304:513-520.
- Dryden, M. S. (2010). Complicated Skin and Soft Tissue Infection. *Journal of Antimicrobial Chemotherapy*, 65, 35-44. <https://doi.org/10.1093/jac/dkq302>.
- Ducel, G., Fabry, J. and Nicolle, L. (2010). Prevention of hospital-acquired infections: A practical guide. *World Health Organization, Department of Communicable Disease, Surveillance and response*, 2nd edition. 10-12.
- Edwards, R. and Harding, K. G. (2004). Bacteria and wound healing. *Journal on Current infectious diseases*, 17: 19-26.
- Egbe, C., Omoregie, R., Igbarumah, I. and Onemu, S. (2011). "Microbiology of wound infections among patients of a Tertiary Hospital in Benin City, Nigeria". *Journal Resource of Health Sciences*, 11(2): 109-113.
- Eliya-Masamba, Martha C. Banda., and Grace W. (22 October 2013). "Primary closure versus delayed closure for non-bite traumatic wounds within 24 hours post injury". *The Cochrane Database of Systematic Reviews*. (10): CD008574.
- Emele, F. E., Izomoh, M. I. and Alufohai, E. (1999). Microorganism Associated with Wound Infection in Ekpoma, Nigeria. *West African Journal in Medicine* 18(2):92-100.PMID:10504864.
- Fonder, M. A., Lazarus, G. S., Cowan, D. A., Aroscook, B., Kohli, A. R. and Mamelak, A. J. (2008). Treating the chronic wounds: A practical approach to the care of non-healing wounds and wound care dressings. *Journal of Academy Dermatology*, 58 (2): 185-206.
- Frykberg, R. G. and Banks, J. (2015). Challenges in the treatment of chronic wounds. *Advanced Wound Care*. 4(9):560-582.
- Goldstein, E. J., Citron, D. M. and Nesbit, C. A. (1996). Diabetic foot infections. *Bacteriology and activity of 10 oral antimicrobial agents against bacteria isolates from consecutive cases*. *Diabetic Care*, 19: 638-641.
- Gould D. (2011). Diagnosis, prevention and treatment of fungal infections. *Nursing Standard*. 20-26;25(33):38-47.
- Gould, L., Abadir, P. and Brem H. (2018). Chronic wound repair and healing in older adults: current status and future research. *Wound Repair Regeneration*. 23(1):1-13.
- Gwaltney, J. M., Moskalski Jr, P. B. and Hendley, J. O. (1978). Hand-to-hand transmission of rhinovirus colds. *Annual International Journal in Medicine*. 88:463-467.
- Health New organization of infected wounds. (2018). Wound care center.

- <https://www.woundcarecenters.org/woundtypes/infected-wounds>.
- Heo, S. H., Shin, S. S., Kim, J. W., Lim, H. S., Seon, H. J. and Jung, S. I. (2014). Imaging of actinomycosis in various organs: a comprehensive review. *Radiographics*. 34 (1):19-33. [Medline].
- Holloway, S. and Jones.V. (2017). The importance of skin care and assessment. *Britian Journal for Nursing*. 14(22):1172–1176.
- International Consensus Update (ICU) (2016). Wound Infection in Clinical Practice: Principles of Best Practice.https://www.woundinfection-institute.com/wp-content/uploads/2017/07/IWII_Consensus_Final-2017.pdf.
- Jarbrink, K., Ni G. and Sonnergren, H. (2018). Prevalence and incidence of chronic wounds and related complications: a protocol for a systematic review. *Systemic Reviews*. 5(1):152.
- Janet, M., Torpy, M. D., Alison, M. A., Richard, M. and Glass, M. D. (2005). Wound infection. *Journal of the American Medical Association*, 294: (16) 2122.
- Jess, S. (2020). Barriers to Infection. Teach me physiology limited. [CC BY-SA 4.0 (<http://creativecommons.org/licenses/by-sa/4.0>)], via Wikimedia Commons.
- Jones and Bartletts, (2011). *American Academy of Pediatrics*. First Aid for Females (39).
- Jull, A. B., Cullum, N., Dumville, J. C., Westby, M. J., Deshpande, S. and Walker, N. (2015). "Honey as a topical treatment for wounds". *Cochrane Database of Systematic Reviews*. 3 (3): CD005083. Könönen E, Wade WG. (2015) Actinomyces and related organisms in human infections. *Clinical Microbiology Revised*. (2):419-442. [Medline].
- Leaper, D., Assadian, O. and Edmiston C. E. (2018). Approach to chronic wound infections. *Brand Journal on Dermatology*. 173(2):351–8.
- Mani R, Margolis D. J and Shukla V. (2018). Optimizing technology use for chronic lower-extremity wound healing: a consensus document. *International Journal on Low Extreme Wounds*. 15(2): 102–119.
- Mahmood, A. (2000). Bacteriology of surgical site infections and antibiotic susceptibility pattern of the isolates at a tertiary care hospital in Karachi. *Journal Pakistan Medical Association*, 50: 256-259.
- Mangram, A. J., Hosan, T. C., Pearson, M. L., Silver, L. C. and Jarvis, W. R. (1999). Guideline for prevention of surgical site infection. Centers for disease control and prevention (CDC). Hospital infection control practices advisory committee. *American Journal on infectious countries*, 27: 97-132.
- Martin, J. M., Zenilman, J. M. and Lazarus, G. S. (2010). Molecular microbiology: new dimensions for cutaneous biology and wound healing. *Journal investment in Dermatology*, 139 (1): 38-48.
- Matsuura, G. T. and Barg, N. (2013). Update on the Antimicrobial Management of Foot Infections in Patients with Diabetes. *Clinical Diabetes*, 31, 59-65.<https://doi.org/10.2337/diaclin.31.2.59>.
- McCulloch, J. M. and Kloth, L. C. (2010). Wound healing Evidence based management. 4th Edition Philadelphia, PA: F.A. Davis company, 2010:(5)109.
- Mulu, W., Kibru, G., Beyene, G. and Damtie, M. (2012). Postoperative nosocomial infections and antimicrobial resistance pattern of bacteria isolates among patients admitted at Felege Hiwot Referral Hospital, Bahirdar Ethiopia. *Ethiopian Journal of Health Science*, 22(1): 7-18.
- Murphree, R. W. (2017). Impairments in skin integrity. *Nursing Journal Clinical North American*. 52(3):405–417.
- National Pressure Ulcer Advisory Panel (NPUAP) (2018). Shear force slide set. <https://www.npuap.org/resources/educational-and-clinical-resources/shear>.
- Nita, P., Nikita, S., Rajni, S., Saroj, H. and Rakesh, K. M. (2018). Prevalence of Multidrug (MDR) and Extensively Drug Resistant (XDR) *Proteus* species in a Tertiary-Care Hospital, India.

- International Journal of Current Microbiology and Applied Sciences*, 3, 243-252.
- Nuutila, C. R. K. and Lee, C. C. Y. (2018). The external microenvironment of healing skin wounds. *Wound Repair Regeneration* .23(4):456–464.
- Obritsch, M. D., Fish, D. N., MacLaren, R and Jung, R. (2004). National surveillance of antimicrobial resistance in *Pseudomonas aeruginosa* isolates obtained from intensive care unit patients from 1993 to 2002. *Antimicrobial Agents of Chemotherapy*, 48:4606-4610.
- O'Dell, M. L. (2018). Skin and wound infection: An overview: *American family physician Journal*, 57(10) 2424-2432.
- Ohalete, C. N., Obi, R. K. and EmeaKoroha, M. C. (2012). Bacteriology of Different Wound Infection and Their Antimicrobial Susceptibility Patterns in Imo State, Nigeria. *World Journal of Pharmaceutical Sciences*, 13, 1155-1172.
- Olayinka, A. T., Onile, B. A. and Olayinka, B. O. (2004). Prevalence of Multi-Drug Resistant (MDR) *Pseudomonas aeruginosa* Isolates in Surgical Units of Ahmadu Bello University Teaching Hospital, Zaria, Nigeria: An Indication for Effective Control Measures. *Annals of African Medicine*, (1): 13-16.
- O'Mathúna, D. P. (2016). "Therapeutic touch for healing acute wounds" (PDF). *The Cochrane Database of Systematic Reviews*. (8): CD002766.
- Omole, A. and Stephen, E. (2014). Antibigram Profile of Bacteria Isolated from Wound Infection of Patients in Three Hospitals in Anyigba, Kogi State, Nigeria. *FUTA Journal of Research in Sciences*, 2, 258-266.
- Pondei, K., Fente, B. G. and Oladapo, O. (2013). "Current microbial Isolates from wound swabs, their culture and sensitivity pattern at the Niger Delta University Teaching Hospital, Okolobiri, Nigeria". *Tropical Medicine and Health*, 41(2): 49-53.
- Rodero, M. P. and Khosrotehrani, K. (2010). Skin wound healing modulation by macrophages. *International Journal of China Experimental Pathology*, 3 (7): 643-653.
- Ron walls, M. D., John, J., Ratey, M. D., Robert, J. and Simon, M. D. (2009). Rosen's emergency medicine: expert consult premium edition. *Concept and clinical practice*, 2: 2482-2483.
- Saeedi, M., Eslamifar, M. and Khezri, K. (2019). Kojic acid applications in cosmetic and pharmaceutical preparations. *Pharmacotherapy* 2019, 110, 582–593.
- Scalise, A., Calamita, R. and Tartaglione, C. (2018). Improving wound healing and preventing surgical site complications of closed surgical incisions: a possible role of incisional negative pressure wound therapy. *A systematic review of the literature. International Wound Journal* 13(6):1260–1281.
- Sahu, S., Shergill, J., Sachan, P. and Gupta, P. (2011). Superficial Incisional Surgical-Site Infection in Elective Abdominal Surgeries. A Prospective Study. *The Internet Journal of Surgery*, 26, 514-524. <https://doi.org/10.5580/14a8>.
- Sandu, C. D., Constanin, C., Raducu, L., Moraru, O. and Visan, S. A. (2018). Management of necrotic lesions in chronic limb ischemia. *Journal on Surgery Science*. 2(1):34–70.
- Sani, R. A., Garba, S. A. and Oyewole, O. A. (2012). Antibiotic Resistance Profile of Gram-Negative Bacteria Isolated from Surgical Wounds in Minna, Bida, Kontagora and Suleja Areas of Niger State. *American Journal of Medicine and Medical Sciences*, 2, 20-24. <https://doi.org/10.5923/j.ajmms.20120201.05>.
- Schnuriger, B., Inaba, K., Eberle, B. M., Wu T, Talving P. and Bukur, M. (2010). Microbiological profile and antimicrobial susceptibility in surgical site infections following hollow viscus injury, *Journal on Gastrointestinal surgery*, 14: 1304-1310.
- Siddiqui, A. R., and Bernstein, J. M. (2010). Chronic wound infection: facts and controversies. *Clinical Dermatology*, 28: 516-526.

- Simple wound management (SWM) (2015). Archived at the Wayback Machine, patient.info (website).
- Stewart, M. G. (2005). Principle of ballistic and penetrating trauma. *Comprehensive Management*, 188-194.
- Sule, A., Thanni, L., Sule-Odu, O. and Olusanya, O. (2002). "Bacterial pathogens associated with infected wounds in Ogun state University Teaching Hospital, Sagamu, Nigeria". *African Journal of Clinical Experimental Microbiology*, 3(1): 13–16.
- Sullivan, D. C. and Chapman, S.W. (2010). Bacteria that masquerade as fungi: actinomycosis/nocardia. 7 (3):216-21. [Medline].
- Tissue. (2018). Severe edema: a detriment to wound healing. Available at: <https://www.advancedtissue.com/severe-edema-a-detriment-to-wound-healing/>.
- Tom, I. M., Agbo, E. B., Umar, A. F., Muhammad, M. I., Askira, M. U., Jidda, B. U., Abdullahi, A. and Ali, B. H. (2018). Plasmid Profile Analysis of Multi-Drug Resistant *Proteus* spp Isolated from Patients with Wound Infection in Northeastern Nigeria. *International Journal of Pathogen Research*, (1):1-9.
- Tomislav Meštrović. (2018). Viral skin infection <https://www.health.auckland.ac.nz/courses/dermatology/2-infections/viral.html>. CD102566.
- Toon, Clare D., Sinha, Sidhartha., Davidson, Brian R., Gurusamy and Kurinchi Selvan (2015). "Early versus delayed post-operative bathing or showering to prevent wound complications". *The Cochrane Database of Systematic Reviews* (7): CD010075.
- Tyring, S. K. (2010). Cutaneous Virology. In: Tyring S. K., Moore A. Y., Lupi O., editors. *Mucocutaneous Manifestations of Viral Diseases, Second Edition*. Information Healthcare, London, UK; pp. 1-19.
- World Union of Wound Healing Societies (WUWHS) (2018). Principles of best practices: wound infection in clinical practice. *An international consensus update*, 2016. London: MEP LIMITED.
- Wound Care Center (WCC). (2020). <https://woundcarecenters.org> HSUBL, 4 cba internet brand.
- Wound Source Practice Accelerator Series (2018). RED SKIN mnemonic for assessment, prevention, and treatment of pressure injuries and ulcers. *WoundSource.com*. <https://pages.woundsource.com/using-the-red-skin-mnemonic-pressure-injuries>. Accessed December 30, 2018.
- Yamada, R., Yoshie, T., Wakai, S., Asai-Nakashima, N., Okazaki, F., Ogino, C., Hisada, H., Tsutsumi, H., Hata, Y., and Kondo, A. (2014). *Aspergillusoryzae*-based cell factory for direct kojic acid production from cellulose. *Microbial Cell Fact.* 13-17.