
Heavy Metals Sludge from Septic Tank Sewage: Implication for Use as Fertilizer

Oyem I. M.^{1*} and Oyem H. H.²

1. Department of Biological Sciences University of Delta, PMB 2090, Agbor Delta State, Nigeria.

2. Department of Chemical Sciences, University of Delta, PMB 2090, Agbor Delta State, Nigeria.

* Corresponding author: mirian.oyem@unidel.edu.ng

Abstract: Sludge contains nutrients, heavy metals, and metalloids some of which are essential macronutrients, such as N, P, K, Ca, Mn and other essential trace elements, such as Cu, Fe and Zn. This paper examines the heavy metals within the septic tank system and their eventual accumulation as components of sludge with implication for agricultural use. Sludge samples from the inlet and outlet chambers of the septic tank were randomly collected from three study locations in the south-south region of Nigeria. The composite samples were made for each of these study locations from which nine trace metals (Fe, Zn, Mn, Pb, Cu, Cr, Cd, Ni and Va) were analysed. The concentrations of the metals were determined using standard methods for water and wastewater examination. Results showed that sludge samples from the anaerobic bioreactors contained a high concentration of various trace metals. Iron had the highest mean concentration level of 556 ± 0.2 mg/kg – 3391 ± 1.13 in the raw sewage sludge sample, while vanadium was the least in abundance with mean concentrations of 0.86, 0.77, and 1.52 mg/kg in locations A, B and C respectively in the inlet chamber sludge samples. Although, sludge is variously used for agricultural purpose, there is need for treatment to reduce the concentration levels of heavy metals prior to its use as fertilizer as a matter of public health importance.

Key word: Fertilizer, food chain, heavy metals, septic tank, sewage sludge

INTRODUCTION

Hheavy metals are elements whose densities are high typically above 5 g/cm³ (Hawkes, 1997; Chu, 2018; Ali *et al.*, 2019; Briffa *et al.*, 2020; Sayo *et al.*, 2020). Examples of heavy metal elements include lead, zinc, chromium, cadmium, arsenic, mercury, vanadium, manganese, and iron. These metals are found in the environment from geogenic and anthropogenic sources (Tchounwou *et al.*, 2012; Ibrahim and Ibrahim, 2016). The metals exist in the air, surface and groundwater, and soil. Thus, can be transported through the soil to plants and up the food chain to man (Nkwunonwo *et al.*, 2020), as well as through air and water pollution. Man is the major receiver of these metals in the environment from the uptake of these heavy metals through plants, surface and groundwater, and airborne pollutants (He *et al.*, 2005; Usman *et al.*, 2020). This makes man a veritable bio-accumulator of these heavy metals which eventually arrive at the septic tank.

Septic tanks are digestion chambers where anaerobic bacteria breakdown organic matter from human domestic activities is treated in

a two-step process effected by acid- and methane-forming microorganisms (Abatenh *et al.*, 2017). The process of microbial remediation of heavy metals in the sewage of septic tank systems is dependent on the pH of the system (Ademoroti, 1996; Abatenh *et al.*, 2017). This therefore, determines the threshold and balance of equilibrium between acid-loving bacteria (acidophiles) and bacteria that thrive in alkaline environment (methanogens). Bio-precipitation of trace metals onto cell surfaces by microbes is favoured by acidic conditions (Ademoroti, 1996; Canovas *et al.*, 2003; Jin *et al.*, 2004; Yuncu *et al.*, 2006; Hassan *et al.*, 2009; Igiri *et al.*, 2018; Osman *et al.*, 2019; Rose *et al.*, 2019; Delangiz *et al.*, 2020;). By convention, septic tank systems are designed to have two chambers; that is, the raw (inlet) chamber and the semi-treated (outlet) chamber. Hence, the components of the septic tank be classified into suspended solids, effluent and sludge.

For septic tank to work optimally, it should be evacuated every 3 to 5 years, however, inspection should be conducted biannually to confirm that the system is working optimally, and that there are no leaks. It is

also necessary to check that the levels of sludge and scum in the tank has not risen drastically (USEPA, 1997) that would necessitate evacuation should it exceed 25 percent of the working liquid capacity of the tank or the level of the layer of scum is within three inches of the bottom of the outlet baffle (USEPA, 1994). These situations make the sludge readily available for utilization for agricultural purpose as organic fertilizers.

Sludge is the visco-solid component of sewage which is about 57.0 per cent water (Ademoroti, 1996). The precipitation of both the solvated and insoluble substances leads to the development of sludge (Diep *et al.*, 2018; Zhang *et al.*, 2021). Sludge is composed of proteins, carbohydrates, pathogens and heavy metals (Zhang *et al.*, 2021). Even though sewage sludge contain a deluge of plant nutrients, the heavy metals content can be a problem (Oudeh, 2002). Therefore, there is a need to treat accumulated sludge in a wastewater before safe disposal especially into water bodies, which is usually the practice. The essence of treating sludge is to reduce the amount of organic matter, potentially toxic heavy metals and the population of disease-causing microorganisms. The most common treatment options include anaerobic digestion. Others, include aerobic digestion and composting. Treated sludge can be disposed off or utilized as fertilizers for agricultural land, providing cheap manure for farmers (El sokkarry, 1993). Waste water sludge is a useful source of nitrogen, phosphorus and organic matter. However, there are limits to what can be applied. For example application of waste water sludge should not exceed 250 kg nitrogen per hectare per annum, particular within nitrate vulnerable zones (EEC, 1986; USEPA, 1997).

Fertilizers are substances which are formulated to aid plant growth, where the normal soil may lack sufficient nutrient to support plants growth and development. Application of inorganic fertilizer to crops has become difficult owing to its expensive

nature and sometimes unavailability. Sludge therefore, provides cheap and readily available means of supplying essential nutrients to plants – Green Science. However, as beneficial as the use of sludge as fertilizer is, the process is frost with some misgivings as waste water sludge contains large concentration of heavy metal than that present in most soils (Gomez-Canela *et al.*, 2012). Once applied, the heavy metals will accumulate in the top soil until it is picked up by plants.

However, the case is being made by green environmentalists for organic fertilizers to make up the heavier source of fertilizers use by farmers. Reasons given for this include, the advantage organic fertilizers have in terms of their composition apart from it having a natural source. Manure as organic fertilizers as often called, possess minerals, organic matter, moisture content, and microbes, some of which could be beneficial to the soil and plant eventually, unlike the inorganic fertilizers whose components are hewn out of source rocks and lack the same compositions as organic fertilizers (Zhou *et al.*, 2022). In spite of the fact that during the formation of manures by microbes, methane gas (a greenhouse gas) is formed which adds to the burden of global greenhouse atmospheric load, manure from septic tanks – a pH controlled anaerobic digester can mitigate this problem (Oyem *et al.*, 2022), while providing plant with the requisite nutrients via sludge. This paper focused on ascertaining the levels of heavy metals present in septic tank sludge and the implications of using sludge as fertilizers.

MATERIALS AND METHODS

Sample area: Sludge samples for analysis were collected from different septic tanks in three locations in Agbor in Delta State, Nigeria. Agbor has a population of 162,594 (National Bureau of Statistics, 1977), with an area covering 436 sq. kilometres.

Sample collection: Sludge samples from septic tank sewage were collected from three locations in Agbor in 1 L plastic bottles. Samples were immediately taken to the

laboratory in cool ice packs. Samples were then made into a composite sample for analyses using the APHA (1995) method of wastewater analysis.

Heavy metal analyses: Heavy metals ions present in the sample were analysed using Atomic Absorption Spectrometry (AAS) instrument, model EDX 800, manufactured in Japan. The source of radiation used was a Hollow lamp which emitted the wavelength characteristics of the metal, using a different lamp source for each metal and directed through a flame into a monochromator that selected the preferred wavelength. A photomultiplier tube was the detector and converted the incident signal into an electric signal. The metals were identified at varying wavelengths and current. In all, nine (9) heavy metal ions were analysed and values obtained were recorded.

RESULTS AND DISCUSSION

Iron recorded the highest concentration of 1410 mg/kg and 615 mg/kg in both the inlet and outlet chambers of the septic tanks sampled in the immediate locality. This would be expected for a region known for its high Fe content from its geology (Avwunudiogba, 2000). The soil is characteristically reddish brown in colour and clearly depicts high iron content. One would therefore expect a high Fe content in this environment, especially in the plants cultivated on this soil and its groundwater. It is therefore not surprising that the difference between the Fe content and other heavy metal ions is significantly high.

Zinc had the second highest heavy metal content of the nine analysed in the sludge component of the study area after iron. Zinc is a trace metal needed in the body's biochemistry, it is necessary for certain important functions of the body (Roohani et al., 2013). Its presence in septic tank to this extent may not be associated entirely to diet and supplements but, also through groundwater (Oyem et al., 2015) from ancient corrugated zinc roofing sheets as well as zinc oxide in most ointments, creams, lotions, shampoos and antiseptics

(NCBI, 2017). Decreased levels of Zn in the semi-treated sludge samples compared to the raw sewage samples was observed. These findings are in tandem with reports by Duan et al. (2017); Steinhardt and Egler (2018) and Agoro et al. (2020), who noted that sludge was a rich deposit of toxic metals and though were within normal range yet in agricultural soils may be a potential risk for human safety.

Manganese concentration in sludge sample analysed was 45.8 mg/kg in the inlet chamber and 17.3 mg/kg in the outlet chamber. These values were the third highest in the study's findings, translating to a significant proportion of the heavy metal load in the environment, a similar pattern was observed as in Zinc.

The sludge sample in the inlet chambers had Cu levels of 13.2 mg/kg and 5.38 mg/kg in the outlet chamber. The Cu is a component of some enzymes in the body besides being ubiquitous as component of many electrical appliances (Mydy et al., 2021). Although, naturally a trace metal, its presence in sludge to this concentration is telling and significant.

Chromium in this study represented as total chromium had a slightly significant value of 1.50 mg/kg and 1.20 mg/kg in both chambers of the septic tanks respectively. Chromium in the environment is treated with concern because of its deleterious effect on health (Oyem et al., 2014; Georgaki and Charalambous, 2022). Although, the values recorded in this study is benign, it is nonetheless considerable.

The concentration of cadmium in the study for the inlet and outlet chambers were 0.02 mg/kg and 0.01 mg/kg. Cadmium presence is often associated with man's industrial activities, and so in areas where Cd concentration is significant, anthropological sources are readily adduced (Singh and Liu, 2024). In this study, Cd concentration is not significant, and this signifies non-industrial activities in the area.

In the inlet chambers' sludge samples ranged from 0.35 to 0.90 mg/kg in the septic tank sewage in the area. Again, these values are

low and do not depict pollution of the area arising from industrial activities (Jakubus and Graczyk, 2020).

In sludge sample Pb concentrations were slightly significant but, not suggestive of serious pollution. The Pb toxicity is treated with serious concern because of its effect on humans. The Pb causes a damage to the central nervous system (WHO, 2023). Therefore, Pb values of 0.90 and 0.50 mg/kg in the inlet and outlet chambers is of slight concern.

Vanadium was the least abundant heavy metal analysed in this study. Vanadium is present in the inlet sludge samples in concentrations of 0.50 mg/kg and 0.30 mg/kg in the study. Vanadium in this area can be traced to the activities of auto mechanics which are present in the area (Wnuk, 2023).

The main risk related to agricultural use of sewage sludge are the potential presence of heavy metals, pathogens and pollutants enrichment in soil, plants and animal

pastures and the subsequent entry into the food chain (Ternes *et al.*, 2004 and Gomez-Canela *et al.*, 2012).

Heavy metals, pathogens and organic pollutants can also affect soil functioning and biodiversity (Harrison *et al.*, 2006 and Roig *et al.*, 2012). According to literature, low application doses of sludge did not cause a significant increase in the heavy metal concentration in soil (Singh and Agrawal, 2008). On the contrary, low metal sludge has beneficial effect on microbial biomass, organic carbon and on soil microbial activity (Usman *et al.*, 2012). Excessive application of sewage sludge to soil has been found to increase the bioavailability of heavy metals that have a negative effect on soil (Singh and Agrawal, 2008 and Usman *et al.*, 2012). However, the option of heavy metal removal using genetically engineered microorganisms has been explored in a recent study (Diep *et al.*, 2018).

Table 1: Heavy metals concentrations (mg/kg) in both chambers of the septic tanks in the study locations

S/N	Metal ion (Analyte)	Inlet chamber (mg/kg)	Outlet chamber (mg/kg)
1.	Fe	1410 ±0.20	615 ±2.00
2.	Zn	110 ±0.20	66.2 ±0.20
3.	Mn	45.8 ±0.20	17.3±0.10
4.	Cu	13.2 ±0.20	5.38 ±0.20
5.	Cr	1.50 ±0.02	1.20 ±0.02
6.	Cd	0.02 ±0.01	0.01±0.05
7.	Ni	0.90 ±0.20	0.35 ±0.01
8.	Pb	0.70 ±0.01	0.50 ±0.02
9.	V	0.50 ±0.02	0.30 ±0.02

CONCLUSION

Heavy metals distribution in the sludge components of septic tank systems was studied to understand their composition within the system. Results showed that sludge samples from the anaerobic bioreactor contained a high concentration of various trace metals. This has high implications for the use of sludge as compost material in Agriculture. Although

sewage sludge is a good source of nutrients for plant growth, the presence of heavy metals in sludge can reduce its use. However, the heavy metals found in this study are not those with great health implications. That notwithstanding, the need to ensure proper treatment of sludge prior to its use as fertilizer is a matter of public health importance.

REFERENCES

- Abatenh, E., Gizaw, B., Tsegaye, Z., and Misganaw, W. (2017). The Role of Microorganisms in Bioremediation - A Review. *Open Journal of Environmental Biology* 2(1): 38-46.
- Ademoroti, C. M. A. (1996a). *Standard methods for water and effluents analysis* (First). Ibadan:
- Ademoroti, C. M. A. (1996b). *Environmental Chemistry and Toxicology* (First). Ibadan: Foludex Press.
- Agoro, M. A., Adeniji, A. O., Adefisoye, M. A., and Okoh, O. O. (2020). Heavy metals in wastewater and sewage sludge from selected municipal treatment plants in eastern Cape Province, South Africa. *Water (Switzerland)*, 12(10): 1–19.
- Ali, H., Khan, E., and Ilahi, I. (2019). Environmental Chemistry and Ecotoxicology of Hazardous Heavy Metals: Environmental Persistence, Toxicity, and Bioaccumulation. *Journal of Chemistry*. 1: 1–14.
- APHA. (1995). *Standard Methods for the Examination of Water and Waste Water* (20th ed.). Washington, DC: American Public Health Association.
- Briffa, J., Sinagra, E., and Blundell, R. (2020). Heliyon Heavy metal pollution in the environment and their toxicological effects on humans. *Heliyon*. 6: 1-26.
- Canovas, D., Cases, I., and de Lorenzo, V. (2003). Heavy metal tolerance and metal homeostasis in *Pseudomonas putida* as revealed by complete genome analysis. *Environmental Microbiology*, 5:1242–1256.
- Chu, D. (2018). Effects of heavy metals on soil microbial community. In *IOP Conference Series: Earth and Environmental Science* 113:1–6.
- Delangiz, N., Varjovi, M. B., Lajayer, B. A., and Ghorbanpour, M. (2020). Beneficial microorganisms in the remediation of heavy metals. In *Molecular Aspects of Plant Beneficial Microbes in Agriculture*. Pp. 417–423.
- Diep, P., Mahadevan, R., and Yakunin, A. F. (2018). Heavy Metal Removal by Bioaccumulation Using Genetically Engineered Microorganisms. *Frontiers in Bioengineering and Biotechnology*. 29. <https://doi.org/https://doi.org/10.3389/fbioe.2018.00157>.
- Duan, B.; Zhang, W.; Zheng, H.; Wu, C.; Zhang, Q. and Bu, Y. (2017). Comparison of Health Risk Assessments of Heavy Metals and As in Sewage Sludge from Wastewater Treatment Plants (WWTPs) for Adults and Children in the Urban District of Taiyuan, China. *International Journal Environmental Research Public Health*. 14:1194.
- El-Sokkary, L.H (1993) Prospects on reuse of sewage waste water and sludge in agricultural land Egyptian Soil Science Society (ESSS) 4th National Congress, Nov, 24-25, Cairo, Egypt.
- European Environmental Commission (1986). EU regulation on waste water management Council Directive 86/278/EEC 12 June. Brussels, Belgium.
- Georgaki, M. and Charalambous M. (2022). Toxic chromium in water and the effect on human body: a systematic review. *Journal of Water and Health* 21(11):1-20.
- Gomez-Canela, C.; Barth, J.A.C. and Lacorte, S. (2012) Occurrence and fate of perflourinated compounds in sewage sludge from Spain and Germany. *Environ. Sci. Pollut. Res.*19: 4109-4119.
- Harrison, E.Z. Oakes; S.R; Hysell, M. and Hay, A. (2006) Organic Chemicals in Sewage Sludge. *Sci. Total Environ*. 367: 481-497.
- Hassan, M. A., Salleh, A., and Milow, P. (2009). Characterization of the Adsorption of the Lead (II) by the Nonliving Biomass *Spyrogia neglecta* (Hassall) Kutzing. *American*

- Journal of Biochemistry and Biotechnology*. 5(2): 75–83.
- Hawkes, S. J. (1997). What is “heavy metals”? *Journal of Chemical Education*, 74(11): 1374.
- He, Z., Yang, X., and Stoffella, P. (2005). Trace elements in agroecosystems and impacts on the environment. *J Trace Elem Med Biol*. 19(2–3): 125–140.
- Ibrahim, M. and Ibrahim, N. (2016). The Relations Between Concentration of Iron and the pH Ground Water (Case Study Zulfi Ground Water). *International Journal of Environmental Monitoring and Analysis*. 4(6): 140.
- Igiri B.E., Okoduwa S.I.R., Idoko G.O., Akabuogu E.P., Adeyi A.O., E. I. (2018). Toxicity and bioremediation of heavy metals contaminated ecosystems from tannery wastewater: A review. *Journal of Toxicology*. 1–16. <https://doi.org/2568028>.
- Jakubus, M., and Graczyk, M. (2020). Availability of Nickel in Soil Evaluated by various Chemical Extractants and Plant Accumulators. *Agronomy*, 10(11): 1805.
- Jin, Y., Luan, Y., Ning, Y., and Wang, L. (2018). Effects and mechanisms of microbial remediation of heavy metals in soil: A critical review. *Applied Sciences*. <https://doi.org/10.3390/app8081336>.
- Mydy, L. S., Chigumba, D. N., and Kersten, R. D. (2021). Plant copper metalloenzymes as a prospect for new metabolism involving aromatic compounds. *Frontiers in Plant Science*. 12: 692108.
- National Bureau of Statistics. (1977). Demographic Statistics. In *Alpacas, Sheep, and Men* Pp. 219–228.
- National Centre for Biotechnology Information Pubchem Compound Database (ID) Sept 16, 2017.
- Nkwunonwo, U. C., Odika, P. O. and Onyia, N. I. (2020). A Review of the Health Implications of Heavy Metals in Food Chain in Nigeria. *The Scientific World Journal* 1–11. <https://doi.org/10.1155/2020/6594109>.
- Osman, G. E. H., Abulreesh, H. H., Elbanna, K., Shaaban, M. R., Samreen, and Ahmad, I. (2019). Recent progress in metal-microbe interactions: Prospects in bioremediation. *Journal of Pure and Applied Microbiology*. <https://doi.org/10.22207/JPAM.13.1.02>.
- Oyem, H. H., and Oyem, I. M., and Usese, A., I. (2015). Iron, Manganese, Cadmium, Chromium, Zinc, and Arsenic groundwater contents of Agbor and Owa communities of Nigeria. *SpringerPlus*. 4:104.
- Oyem, H. H., & Oyem, I. M. (2022). Heavy Metals pH-Mediated Microbial-Remediation in Septic Tank Effluents. *International Journal of Environmental Chemistry and Ecotoxicology Research*. 4(1): 15–35.
- Oudeh, M. (2002). Effects of sewage sludge application on growth and mineral composition of maize plant. Congress of recent technologies in agriculture 1: 280-289. Cairo University, Egypt.
- Rose, J., Jiménez Cisneros, B., Adegoke, A. A. and Stenstrom, T. A. (2016). Septic Systems. In J. B. Rose & B. Jiménez-Cisneros (Eds.), *Water and Sanitation for the 21st Century: Health and Microbiological Aspects of Excreta and Wastewater Management (Global Water Pathogen Project)* Michigan: Michigan State University. Pp. 1–16.
- Roig, N; Sierra, J.; Nadal, M.; Marti, E.; Navalon-Madrigal, P.; Schuhmacher, M. and Domingo, J.L. (2012) Relationship between pollutant content and ecotoxicity of sewage sludge from Spanish waste water treatment plants. *Sci. Total Environ*. 425: 99-109.

- Roohani, N., Hurrell, R., Kelishadi, R., and Schulin, R. (2013). Zinc and its importance for human health: An integrative review. *Journal of Research in Medical Science*. 18 (2): 144-157.
- Sayo, S., Kiratu, J. M., & Nyamato, G. S. (2020). Heavy metal concentrations in soil and vegetables irrigated with sewage effluent: A case study of Embu sewage treatment plant, Kenya. *Scientific African*. 8, e00337. <https://doi.org/10.1016/j.sciaf.2020.e00337>.
- Singh, R., and Liu, Y. (2024). Fate and behaviour of cadmium in soils and recent advances in remediation technologies. In: Ravi Naidu Ed., Inorganic contamination and radionuclides. *Elsevier*. 41-67.
- Steinhardt, G. C., Egler, C. J. and Carrol, N. (2018). *Home & Environment: operating and maintaining and onsite sewage system*.
- Singh, R.P.; Agrawal, M. (2008). Potential benefits and risks of land application of sewage sludge. *Waste Manag*. 28: 347-358.
- Ternes, T.; Joss, A.; Siegrist, H. (2004). Scrutinizing pharmaceuticals and personal care products in wastewater treatment. The complexity of these hazards should not be underestimated. *Environ. Sci. Technol*. 38: 392-399.
- Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., & Sutton, D. J. (2012). Molecular, clinical and environmental toxicology volume 3: Environmental Toxicology. *Molecular, Clinical and Environmental Toxicology*. 101:133–164.
- United State Environmental Protection Agency (1994). Guides to Septage treatment and Disposal. EPA 625/R-94/002 Washington DC.
- United State Environmental Protection Agency (1997). Standards for the use of or disposal of sewage sludge. Final rules. Fed. Reg. Washington DC. 58(32): 9248-9415.
- Usman, K., Khan, S., Ghulam, S., Khan, M.U., Khan, N., Khan, M.A. and Khalil, S.K. (2012) Sewage sludge: An important biological resource for sustainable agriculture and its environmental implications. *Am. J. Plant Sci*. 3: 1708-1721.
- Wnuk, E. (2023). Mobility, Bioavailability, and Toxicity of Vanadium Regulated by Physicochemical and Biological Properties of the Soil. *Journal of Soil science and Plant nutrition*. 23: 1386-1396.
- World Health Organization (2023). Lead Poisoning: Fact sheets. Geneva.
- Yuncu, B., Sanin, F. D. and Yetis, U. (2006). An Investigation of Heavy Metal Biosorption in relation to C/N ratio of activated sludge. *Journal of Hazardous Materials*. 137: 990–997.
- Zhang, X., Ye, P., Wu, Y., Guo, Z., Huang, Y., Sun, Y. and Zhang, H. (2021). Research on dewatering ability of municipal sludge under the treatment of coupled acid and microwave. *Hindawi*. 2021: 1–11.
- Zhou, W., Qing, C., Guo, S., He, J., Deng, X., Song, J., and Xu, D. (2021). Labour off-farm employment and farmers cooking clean energy use: Evidence from rural China. *Agriculture*. 12: 972.