

Occurrence of Metallo-beta-lactamase in Multidrug Resistant *Escherichia coli* Isolated from Donkey Slaughter Market

Ugbo E. N.^{1*} Ugbo A. I.² Agumah B. N.¹ Nwojiji E. C.² Ewah C. M.³ Ogbonna I. P.² and Ekpono C. U.²

1. Department of Applied Microbiology, Faculty of Science, Ebonyi State University, P.M.B. 053, Abakaliki, Ebonyi State, Nigeria.
2. Department of Microbiology and Parasitology, Faculty of Allied Health Sciences, David Umahi Federal University of Health Sciences, P. M. B. 211 Uburu, Ebonyi, Nigeria
3. Department of Biochemistry, Faculty of Science, Ebonyi State University, P.M.B. 053, Abakaliki, Ebonyi State, Nigeria

* Corresponding author: emmanuel.ugbo@ebsu.edu.ng; ugbonuel2001@yahoo.com

Abstract: Animals used in food production have been identified as one of the major primary sources of antibiotic resistant pathogenic bacteria within the Enterobacteriaceae family. The *Escherichia coli* harboring metallo- β -lactamases (MBLs) is a serious threat to global health. This study was aimed at the occurrences of metallo-beta-lactamase in multidrug resistance *Escherichia coli* isolated from donkey slaughter market. A total of 75 swab samples were collected from equipment used in donkey slaughters and *E. coli* was identified using Eosin Methylene Blue Agar (EMBA), MacConkey Agar (MCA) media and other microbiological standard techniques. The *E. coli* isolates were tested for multidrug resistance (MDR) using disk diffusion method and MARI was calculated. The presence of metallo-beta-lactamase (MBL) in multidrug resistance *E. coli* was confirmed using modified Hodge test method. Thirty (40.0%) swab samples were positive for *E. coli*, knife had 8(32.0%), table had 12(48.0%) and slab harbored 10(40.0%). The isolates were resistance to tetracycline (40.0%-80.0%), gentamicin (25.0%-30.0%), erythromycin (40.0%-50.0%), ampicillin (70.0%-75.0%) and ciprofloxacin (20.0%-37.5%). Three major antibiotics resistance pattern were revealed from the 14 (46.7%) isolate that were MDR-*E. coli* which includes; TE-CN-E-AMP-CIP, TE-E-AMP-CIP and TE-CN-E-AMP. This study revealed the occurrences of MBLs in MDR- *E. coli* to be 5(35.7%). The presence of MBLs in MDR *E. coli* isolated from donkey creates strong threat to the treatment of such infection in clinical setting and it calls for an urgent veterinary surveillance program to monitor antibiotics used as growth enhancers in animal production.

Key word: Metallo-beta-lactamase, MDR *E. coli*, donkey slaughter equipment, public health

INTRODUCTION

The livestock industry especially donkey plays a vital role in various regions such as in Asia, Middle-east, Africa like Nigeria, where it is a significant source of meat and milk consumption (FAO, 2019). Despite its importance, concerns about food safety and potential zoonotic diseases have been raised some years ago (Mauro *et al.*, 2016). One of the potential risks associated with donkey meat is the presence of pathogens, such as *Escherichia coli* which can lead to gastrointestinal illnesses in humans when the contaminated product is consumed (USDA-FSIS, 2021). The transmission of *E. coli* from animals to humans has been a subject of increasing public health concern, particularly when it involves the food production chain (CDC, 2021). The improper handling and processing of donkey

meat can potentially introduce pathogenic *E. coli* and other bacterial contamination (Wang *et al.*, 2017). The donkey scientifically known as *Equus africanus asinus* is a domesticated member of the horse family, Equidae (Orhan *et al.*, 2012). Donkeys have been used as a work animal for at least 5000 years. They are mostly found in under developed countries where they are used principally as draught or pack animals.

In developing countries donkeys are valued in particular for their ability to survive under harsh conditions (Swai and Bwanga, 2018). Interestingly, donkey faeces is sometimes used in rural communities to rub/coat the inner walls of mud buildings where human beings live, as local fertilizers and this creates a strong potential for contamination and/or infection of those persons who perform this work and people who live in

such houses, touch these surfaces and farmers who use it for crop productions (Pritchard *et al.*, 2019). Furthermore, it has been established that donkeys shed *E. coli* (Jesse *et al.*, 2015) which poses a great risk to the people handling these animals directly or indirectly, it could also expose these people to diseases caused by this organism. In developing countries, including Nigeria, Ethiopia, animals are commonly slaughtered and processed under unhygienic conditions and these further compromises the microbiological quality and safety of the meat obtained from the animals (Bello *et al.*, 2015; Dulo *et al.*, 2015).

Food-producing animals such as donkey harboring multidrug resistance genes together with genes that mediate the production of some high-profile antibiotic hydrolyzing enzymes such as metallo-beta-lactamases (MBLs), extended spectrum beta-lactamases (ESBLs) and AmpC enzymes possess health risks to the human population particularly due to their potential contribution to the spread of multidrug resistant microorganisms in the community (Ejikeugwu *et al.*, 2018). MBL-encoding genes are usually carried by mobile genetic elements that facilitate horizontal gene transfer (HGT) between bacteria and harbor a great ability to spread (Pierce *et al.*, 2017). MBL-producing bacteria are regarded as the most important environmental pathogens, and further spread of them in the healthcare settings will pose a serious global threat in the future. Therefore, active surveillance is needed to detect the prevalence and incidence of MBL-producing bacteria in the environment and help prevent the spread of these organisms (Beresford and Maley, 2019).

The worldwide distribution of *E. coli* harboring metallo- β -lactamases (MBLs) and AmpC β -lactamases (AmpC) is a serious threat, and due to MBL production, carbapenem resistance is progressively spreading among clinical isolates of *E. coli* (Tewari *et al.*, 2018). Treatment of infections has been complicated by the emergence of multidrug-resistant (MDR) strains of *E. coli*

(Aghil *et al.*, 2021). Food-producing animals have been identified as the primary source of antibiotic-resistant pathogenic bacteria within the Enterobacteriaceae family across multiple countries (Dey *et al.*, 2023; Sobur *et al.*, 2019; Ejikeugwu *et al.*, 2016). Among the most encountered members of Enterobacteriaceae in this category are *Escherichia coli*, *Klebsiella* species, *Shigella* species, and *Salmonella* species, owing to their role in causing diseases in livestock (de Souza *et al.*, 2023; Ugbo *et al.*, 2023). Little information is available on the prevalence of MBLs in MDR *Escherichia coli* from donkey. However, investigating metallo-beta-lactamases (MBLs) in this bacterium is of utmost importance to address, manage the emergence and spread of antibiotic-resistant strains among food-producing animals, it is also a strong critical measure for safeguarding both animal and human health. Thus, the need for studying the occurrences of metallo-beta-lactamase in multidrug resistance *Escherichia coli* isolated from donkey slaughter market.

MATERIALS AND METHODS

Sample collection and processing: Exactly 75 samples (25 each from knives, tables and slabs) were taken aseptically from donkey abattoir at Ezzamgbo, Ohaukwu Ebonyi State using sterile swab stick. The sterile swab sticks were used to collect the samples from the donkeys slaughter equipment by rotating at an angle of 180°C. The swab sticks were returned to their respective containers and labeled properly. The samples were immediately transported to the Applied Microbiology Laboratory of Ebonyi State University, Abakaliki for bacteriological analysis. Each of the collected (knives, tables and slabs) samples were inserted into 5 ml of freshly prepared nutrient broth and the tubes were loosely covered with cotton wool. The tubes were arranged on test tube rack and were incubated at 37°C for 18-24 hours. Bacterial growth was suspected by the presence of turbidity or cloudiness in the tubes after incubation. Tubes that showed turbidity

were further sub-cultured onto solid culture media plates for the isolation of the primary bacterium (Awosile *et al.*, 2018; Ejikeugwu *et al.*, 2018).

Isolation and Identification of *Escherichia coli*: The bacterial colonies obtained were further inoculated on Eosin Methylene Blue Agar (EMBA) (HiMedia M317) and MacConkey Agar (MCA) (HiMedia MH081) media and incubated at 37°C for 18-24 hours for the purpose of isolation of *E. coli*. Further identification of *E. coli* was done using Gram staining and other standard microbiological methods which includes the biochemical test; Triple Sugar Iron Agar (TSIA) (HiMedia M021); Simmons Citrate Agar (SCA) (HiMedia M099), IMVIC media, such as Sulfide Indole Motility (SIM) (HiMedia M181); Methyl red (MR); Voges-Proskauer (VP) (Merck; 105712) (Yanestria *et al.*, 2022).

Antibiotic susceptibility testing: Susceptibility testing was done on Mueller Hinton agar plates (Oxoid, UK) using the Kirby-Bauer disk diffusion method as per the criteria of Clinical Laboratory Standard Institute (CLSI, 2020). The different classes of antibiotics disk which includes Fluoroquinolone (CIP) ciprofloxacin; 5µg, Macrolides (E) erythromycin; 30µg, Tetracycline (TE) tetracycline; 30 µg, Beta-lactams (AMP) ampicillins; 30µg, and Aminoglycosides (CN) gentamycin; 500µg. All the antibiotic disks were procured from Oxoid limited (Oxoid, UK). A loopful of the test organism (adjusted to 0.5 McFarland turbidity standards) was streaked on freshly prepared Muller-Hinton agar plates; and the plates were allowed to stand for 15 minutes. The antibiotic disks were placed at a distance of 30mm apart from each other and 15mm away from the edge of the plate and the susceptibility plates were incubated at 37°C for 24 hours (Ugbo *et al.*, 2023; CLSI, 2020). The zones of inhibition diameter were measured according to the CLSI criteria.

Multidrug Resistance Index: The MAR index for a single isolate was calculated as the number of antibiotics to which an isolate

is resistant to (a) divided by the total number of antibiotics tested against the isolate (b) (Ejikeugwu *et al.*, 2017).

Screening for the presence of metallo-β-lactamase (MBL): All the multidrug resistance *Escherichia coli* isolates were screened for the production of MBL by determining their susceptibility to any of the carbapenems including imipenem (IPM) (10 µg), meropenem (MEM) (10 µg), ertapenem (ETP) (10 µg) (Dey *et al.*, 2023). The Kirby-Bauer disk diffusion technique were used, and each of the antibiotics disk were placed at a distance of 20 mm apart and the plates were incubated at 37°C for 18-24hours. MBL enzyme-producing isolates were suspected when the test organism(s) showed reduced susceptibility to any of the tested antibiotics. The isolates showing inhibition zone diameter (IZD) of ≤ 23 mm were suspected to produce MBL and these isolates were subjected to phenotypic confirmation test according to the method of (Ejikeugwu *et al.*, 2016).

Phenotypic detection of MBL: The multidrug resistance *Escherichia coli* isolates found to be resistant to imipenem or meropenem as identified in the screening test were subjected to phenotypic studies for the presence of metallo β-lactamase (MBL). The pure culture of the multidrug resistance *Escherichia coli* isolates were adjusted to 0.5 McFarland turbidity standards and aseptically swabbed on Mueller-Hinton (MH) agar plates. The standard antibiotic disks of imipenem (10 µg) and meropenem (10 µg) impregnated with EDTA (1 µg) were aseptically placed on MH agar plates. Moreso, supplementary imipenem (10 µg) and meropenem (10 µg) disks without EDTA were also placed alongside with the antibiotic disks impregnated with the chelating agent (EDTA) at a distance of 20 mm away from each other. The chelating agents were initially tested on the test bacteria prior to the phenotypic assay to confirm there were no inhibitory effect on the test organisms. All the plates were incubated at 37°C for 18-24 hours and zone of inhibition were recorded after incubation.

A difference of ≥ 7 mm between the zones of inhibition of any of the carbapenem disks with or without the chelating agents infers metallo-beta-lactamase production according to (Bajracharya *et al.*, 2023).

RESULTS

Of the 75 donkey slaughter equipment samples analyzed, only 30 samples were *E. coli* positive with percentage profile of 40.0%. The result revealed that knife harbored *E. coli* 8(32.0%), table had 12(48.0%) and slab harbored 10(40.0%) (Table 1). The *E. coli* isolated from the slaughter equipment presented different kinds of resistance to the tested antibiotics; tetracycline (40.0%-80.0%), gentamicin (25.0%-30.0%), erythromycin (40.0%-

50.0%), ampicillin (70.0%-75.0%) and ciprofloxacin (20.0%-37.5%). However, most of the isolates studied showed high level of susceptibility to ciprofloxacin (80.0%) and gentamicin (75.0%) (Table 2). Out of the 30 *E. coli* isolates identified from donkey slaughters equipment 14 (46.7%) were observed to present multidrug resistance with three different patterns which includes; TE-CN-E-AMP-CIP, TE-E-AMP-CIP and TE-CN-E-AMP (Table 3). Thus, the 14 MDR *E. coli* isolates, knife had 3 (37.5%), table had 6 (50.0%) and slab harbored 5 (50.0%). Five (35.7%) MDR *E. coli* isolates were confirmed to produce metallo β -lactamase (MBLs), knife 1 (33.3%), table 2 (33.3%) and slab 2 (40.0%) (Table 4).

Table 1: Distribution of *E. coli* isolates from donkey slaughter equipment

Sample/Source	No of Samples	No Positive for <i>E. coli</i>	Percentage (%)
Knife	25	8	32.0
Table	25	12	48.0
Slab	25	10	40.0
Total	75	30	40.0

Table 2: Antimicrobial susceptibility profile of *E. coli* isolated from donkey slaughter equipment

SOURCE	TE		CN		E		AMP		CIP	
	S(%)	R(%)	S(%)	R(%)	S(%)	R(%)	S(%)	R(%)	S(%)	R(%)
Knife	4(50.0)	4(50.0)	6(75.0)	2(25.0)	4(50.0)	4(50.0)	2(25.0)	6(75.0)	5(62.5)	3(37.5)
Table	7(58.3)	5(41.7)	9(75.0)	3(25.0)	7(58.3)	5(41.7)	3(25.0)	9(75.0)	8(66.7)	4(33.3)
Slab	2(20.0)	8(80.0)	7(70.0)	3(30.0)	6(60.0)	4(40.0)	3(30.0)	7(70.0)	8(80.0)	2(20.0)

Key: E = erythromycin, CIP = ciprofloxacin, AMP = ampicillin, CN = gentamicin, TE = tetracycline

Table 3: Multidrug resistance patterns of *E. coli* isolates from donkey slaughter equipment

Isolates Code	Isolates	Number of antibiotics that isolates were resistant to (a)	Total number of antibiotics tested (b)	MAR index (a/b)
K 13	<i>E. coli</i>	TE-E-AMP-CIP	5	0.80
K 24	<i>E. coli</i>	TE-CN-E-AMP-CIP	5	1.00
T 4	<i>E. coli</i>	TE-CIP-E-AMP	5	0.80
T 16	<i>E. coli</i>	TE-E-AMP-CN-CIP	5	1.00
T 21	<i>E. coli</i>	TE-E-AMP-CIP	5	0.80
T23	<i>E. coli</i>	TE-CN-E-AMP	5	0.80
S 2	<i>E. coli</i>	TE-E-AMP-CN-CIP	5	1.00
S 9	<i>E. coli</i>	TE-CN-E-AMP-CIP	5	1.00
S 14	<i>E. coli</i>	TE-E-AMP-CIP	5	0.80
Average MDR index				8/9 (0.88)

Key: K = Knife; T = Table; S = Slab; CIP = ciprofloxacin; E = erythromycin; TE = tetracycline; AMP = ampicillin; and CN = gentamicin

Table 4: Prevalence of MDR-*Escherichia coli* producing metallo β -lactamase (MBLs)

Sample/Source	No of <i>E. coli</i> screened	No of MDR- <i>E. coli</i> suspected for MBL (%)	No of MDR- <i>E. coli</i> +ve for MBL (%)	No of MDR- <i>E. coli</i> -ve for MBL (%)
Knife	8	3 (37.5)	1 (33.3)	2 (66.7)
Table	12	6 (50.0)	2 (33.3)	4 (66.7)
Slab	10	5 (50.0)	2 (40.0)	3 (60.0)
Total	30	14 (46.7)	5 (35.7)	9 (64.3)

DISCUSSION

The global emergence and fast spread of pathogenic microorganisms showing multidrug resistance to antimicrobial agents is a serious public health threat. *Enterobacteriaceae* especially *Escherichia coli* producing metallo beta-lactamase (MBLs) is one of the mechanisms associated with severe bacterial infections in clinical setting. Multidrug resistant properties of these microorganisms have limited and complicated treatment options. This study recorded presence of *E. coli* in donkey slaughter equipment with high level of prevalence at percentage of 40.0. The *E. coli* isolates has been reported by previous studies on donkey slaughter equipment, Fasanmi *et al.* (2019) reported high prevalence of *E. coli* in donkeys, slaughter equipment and their environment in Nigeria. Another study revealed a significant prevalence of *E. coli* strains and raising concerns about potential transmission to humans through contaminated slaughter equipment (Awosile *et al.*, 2018). A study by previous researchers identified various virulence factors in the isolated *E. coli* strains, emphasizing the need for comprehensive screening methods to detect pathogenic strains (Sadiq *et al.*, 2020). The detection of *E. coli* on donkey slaughter equipment raises concerns about the potential for contamination of meat products and the transmission of this pathogen to consumers (Adanech and Temesgen, 2018). *E. coli* is a diverse group of bacteria which is a normal flora of in the gastrointestinal tract of animals which are harmless, but can be pathogenic in case of disease condition in the animals. Thus, in the context of food safety, the presence of pathogenic *E. coli* on slaughter equipment is a critical issue that requires attention to prevent food-borne

illnesses (Yanestria *et al.*, 2022; Adanech and Temesgen, 2018). A study by Mauro *et al.* (2016) on the microbial contamination of cattle carcasses and slaughter equipment in a cattle processing plant observed the presence of *E. coli* on both carcasses and equipment, emphasizing the importance of maintaining hygienic practices to prevent contamination. Several other studies have also investigated the presence of *E. coli* in animal such as goat, sheep, cattle, donkey, chickens slaughter environments and reported a great level of prevalence (Adewunmi *et al.*, 2018; Smith *et al.*, 2018; Johnson *et al.*, 2020; Esonu *et al.*, 2022; Ugbo *et al.*, 2023). The identification of *E. coli* on animals slaughter equipment underscores the importance of implementing rigorous hygiene practices and sanitation measures in abattoirs to minimize the risk of contamination.

The *E. coli* isolates recovered from the slaughter equipment presented different degrees of resistance to the tested antibiotic classes; tetracycline (40.0% to 80.0%), gentamicin (25.0% to 30.0%), erythromycin (40.0% to 50.0%), ampicillin (70.0% to 75.0%) and ciprofloxacin (20.0% to 37.5%). High level of resistance to tetracycline, ampicillin as reported in this study is in accordance with the observation of Mahmoodi *et al.* (2020) who reported *E. coli* resistance to tetracycline as (17.5% to 42.3%), ampicillin (24.6% to 64.7%), low level of resistance to ciprofloxacin (5.2% to 25.4%) and gentamicin (1.9% to 6.5%). Ejikeugwu *et al.* (2016) reported similar level of resistance on *E. coli* isolates from abattoir environment, where they observed that ciprofloxacin showed resistance of 28.1% and gentamicin (18.8%). High level of resistance has been reported on *E. coli* isolates by previous researcher to ciprofloxacin (66.1%) (Nazmul *et al.*, 2019).

A study on pathogenic *E. coli* isolates from chicken meat in Bharatpur, Chitwan reported lower resistance level to ciprofloxacin 16.67% and gentamicin (33.3%) (Bajracharya *et al.*, 2023) and is in line the findings of this study which reported resistance to ciprofloxacin to be (20.0%) and gentamicin (25.0%). Out of the thirty *E. coli* isolates identified from donkey slaughters equipment, 14 (46.7%) were recorded to present multidrug resistance. Multidrug resistance *E. coli* (MDR *E. coli*) isolates in this study showed resistance to four and five different classes of antibiotics with three major different patterns which includes; TE-CN-E-AMP-CIP, TE-E-AMP-CIP and TE-CN-E-AMP. Similar findings of antibiotic resistance pattern includes; ATM-CIP-CN-E, ATM-CIP-C-E, ATM-CN-E, ATM-CIP-E, CIP-C-E were reported on multidrug resistance avian pathogenic *E. coli* (MDR-APEC) isolated from ducks on wet markets in Surabaya (Kendek *et al.*, 2024). Multidrug resistance *E. coli* isolates was reported to show resistance to three (CTX, TZP, IMP (19.0%)); four (ATM, MEM, CIP, CN (23.0%)); five (MEM, AMC, FEP, AMP, CEX (15.0%)) and six (AMC, CTX, ATM, IMP, CIP, CEX (5.0%)) different classes of antibiotics (Nazmul *et al.*, 2019). Previous study on multidrug resistance *E. coli* from animal source observed that isolates were resistance to three and five different classes of antibiotic; AMOX, DO, COT (41.61%), AMOX, DO, COT, AZM (25%), and AMOX, DO, COT, AZM, GEN (16.6%) (Bajracharya *et al.*, 2023). The detection of MDR *E. coli* in slaughter equipment showing resistance to three and five different classes of antibiotics suggests that those classes of antibiotics may have be abused as growth promoters, and during treatment of infections in the animals.

This study revealed the occurrences of metallo β -lactamase (MBLs) in MDR- *E. coli* to be 5 (35.7%). Out of the five (35.7%) MDR *E. coli* isolate that were confirmed to produce metallo β -lactamase (MBLs), knife had 1 (33.3%), table harbored 2 (33.3%) and slab had 2 (40.0%). The study done on *E.*

coli isolated from animals which includes cow and cloacae swabs of poultry birds revealed the presence of MBLs phenotypically to be 39.9% and 45.7% respectively (Ejikeugwu *et al.*, 2017) and is in line with the present report of this study. The prevalence of MBL-producing *E. coli* isolates in this study is in accordance with the observation of the occurrence of MBL-producing *E. coli* isolates (Chouchani *et al.*, 2011). Metallo β -lactamase (MBLs) producing *E. coli* (5.0%) was reported from animal (chicken) (Dey *et al.*, 2023), and 66.0% in clinical (Mahmoodi *et al.*, 2020). Another study from animal reported similar prevalence of *E. coli* isolates positive for MBL production in India (Chakraborty *et al.*, 2010). The presence of MBLs in MDR *E. coli* is a significant concern as it has been linked to numerous outbreaks of foodborne illnesses in humans and this observation highlights a potential foodborne public health hazard (Madec *et al.*, 2017) thus, the need for regular screening of MBLs from animal. The emergence of MBL-production in MDR *E. coli* isolates is alarming since they carry mobile genetic elements with great ability to spread in the environment, thus, early detection of these isolates particularly their reservoir, is crucial to prevent their inter- and intra-care setting dissemination and establish suitable antimicrobial therapies (Mahmoodi *et al.*, 2020).

CONCLUSION

Occurrences of MBLs in MDR *E. coli* (35.7%) isolates which is at high prevalence were detected from donkey slaughter equipment, which suggest serious public health threat. The identification of MDR *E. coli* on slaughter equipment underscores the need for stringent hygiene practices in abattoirs to mitigate contamination risks and ensure consumer safety. However, the encouragement of frequent research and accurate detection of MBL- production in MDR *E. coli* isolates from animal source, abattoir environment and clinical samples is of utmost public health importance due to its

multidrug resistant properties harbored by these organisms to safeguarding both animal

and public health.

REFERENCES

- Adanech, B. H. and Temesgen, K. G. (2018). Isolation and identification of *Escherichia coli* and *Edwardsiella tarda* from fish harvested for human consumption from zeway Lake, Ethiopia. *African Journal of Microbiology Research*, 12: 416-480.
- Adewunmi, O. O., Fagbo, S. Awoyomi, O. J. and Agbaje, M. (2018). Antimicrobial susceptibility pattern of *Escherichia coli* isolates from cattle, goats, and sheep in Nigeria. *Journal of Microbiology and Biotechnology*, 12(2): 139-144.
- Aghil, B., Saeed, K., Nader, H., Melika, M., Mohammadmahdi, K., Nahid, J. and Morteza, S. (2021). Identification of metallo- β -lactamases and AmpC production among *Escherichia coli* strains isolated from hemodialysis patients with urinary tract infection. *Molecular Biology Reports*, 48:7883–7892.
- Awosile, B.B., Oranusi, S.U., Oyekunle, M.A., Dipeolu, M.A., Fagbohun I.K., Adetosoye A.I. and Ojo O.E. (2018). Prevalence and antimicrobial resistance profiles of *Escherichia coli* isolated from donkeys in Nigeria. *Vet Ital.* 54(3): 197-204.
- Bajracharya, A. M., Kharel, P., Thapa, S. and Shrestha, S. (2023). Metallo- β -Lactamase Producing Gram-Negative Bacteria Isolated from Chicken Meat in Bharatpur, Chitwan. *Med. Sci. Forum*, 24: 1.
- Bello, M., Lawan, M. K., Aluwong, T. and Sanusi, M. (2015). Management of slaughter houses in northern Nigeria and the safety of meat produced for human consumption. *Food Control*, 49: 34–39.
- Beresford, R. W. and Maley, M. (2019). Reduced incubation time of the modified carbapenem inactivation test and performance of carbapenem inactivation in a set of carbapenemase-producing Enterobacteriaceae with a high proportion of blaIMP isolates. *J Clin Microbiol.*, 57(7):e01852–18.
- CDC (2021). Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Division of Foodborne, Waterborne, and Environmental Diseases (DFWED). Report on food borne illness.
- CLSI (Clinical Laboratory Standard Institute) (2020). Performance standard for antibiotics susceptibility testing. 11th information supplement NCCLS Documents M100-991 (ISBN 1-56238426-0) NCCL, Pennsylvania, USA.
- Chakraborty, D., Basu, S. and Das, S. (2010). A study on infections caused by metallo-beta-lactamase producing gram-negative bacteria in intensive care unit patients. *American Journal of infectious Diseases*, 6:34-39.
- Chouchani, C., Marrakchi, R., Ferchichi, L., El-Salabi, A. and Walsh, T. R. (2011). VIM and IMP metallo- β -lactamases and other extended-spectrum β -lactamases in *Escherichia coli* and *Klebsiella pneumoniae* from environmental samples in a Tunisian hospital. *APMIS*, 119:725-732.
- de Souza, Z. N., de Moura, D. F., Campos, L. A. A., Córdula, C. R. and Cavalcanti, I.M.F. (2023). Antibiotic Resistance Profiles on Pathogenic Bacteria in the Brazilian Environments. *Arch. Microbiol.*, 205: 185.
- Dey, T. K., Lindahl, J. F., Lundkvist, Å., Grace, D., Deka, R. P., Shome, R., Bandyopadhyay, S., Goyal, N. K., Sharma, G. and Shome, B.R. (2023).

- Analyses of Extended- Spectrum-beta Lactamase, Metallo-Lactamase, and AmpC- Bete Lactamase Producing Enterobacteriaceae from the Dairy Value Chain in India. *Antibiotics*, 12: 1449.
- Dulo, F., Feleke, A., Szonyi, B., Fries, R., Baumann, M. P. O and Grace, D. (2015). Isolation of Multidrug-Resistant *Escherichia coli* O157 from Goats in the Somali Region of Ethiopia: A Cross-Sectional, Abattoir-Based Study. *Plos one*, 10(11): e0142905. doi:10.1371/journal.pone.0142905
- Ejikeugwu, C., Afiukwa, N., Nwuzo, A., Ovia, K., Adonu, C., Ujam, T., Nworie, C., Agbai, E. N., Nwankwo, C. and Onwuenwunor, B. O. (2018). Phenotypic detection of metallo- β -lactamase (MBL) and AmpC enzymes among abattoir isolates of *Escherichia coli* and *Klebsiella* species in Abakaliki, Nigeria. *Trends Med*, 18:1-4.
- Ejikeugwu, C., Esimone, C., Iroha, I., Okonkwo, E. C., Gugu, T., Oli, A. N., Ugwu, M., Ezeador, C. and Moses, N. I. (2017). Detection of Metallo- β -lactamase (MBL) among Carbapenem-resistant Gram-negative Bacteria from Rectal Swabs of Cow and Cloacae Swabs of Poultry Birds. *Ann Med Health Sci Res.*, 7: 51-56.
- Ejikeugwu, C., Edeh, C., Iroha, I., Orji, J., Eluu, S., Ugbo, E., Duru, C. and Esimone, C. (2016). Antibigram and Detection of Metallo-Beta-Lactamase (MBL) positive *Escherichia coli* isolates from abattoir. *Nature and Science*, 14(11): 65-69.
- Esonu, M. C., Kwanashie, C. N, Mamman, P. H. and Esonu, D. O. (2022). characterization and antibiotic susceptibility of *Escherichia coli* isolated from selected internal organs of donkeys slaughtered in abattoirs in Kaduna State, Nigeria. *Nig. Vet. J.*, 43 (2): 1 - 11.
- Fasanmi, A. O., Sulaimon, M. A. and Olufemi, A. S. (2019). Prevalence and Antimicrobial Resistance Profile of *Escherichia coli* Isolated from Donkeys Slaughtered at Ikare Abattoir, Ondo State, Nigeria. *Vet Med Int.*, 3457092.
- Food and Agriculture Organization (FAO) and World Health Organization (WHO). (2019). Microbiological Risk Assessment Series No. 5: Risk Assessment of Shiga Toxin-Producing *Escherichia coli* in Beef and Beef Products. Available from: <http://www.fao.org/3/ca2546en/CA2546EN.pdf>
- Jesse, T. J., Aminu, S., Faruk, M. T., Mikail, B. A., Muhammed, K. G., Jacob, P. K. and Folorunsho, F. (2015). Predictors and risk factor for the intestinal shedding of *Escherichia coli* O157 among working donkeys (*Equus asinus*) in Nigeria. *Veterinary record open*, 2(1): 456-457.
- Johnson D., Williams E. and Brown K. (2020). Genetic Diversity of *Escherichia coli* Strains Isolated from Animal Slaughter Settings: Implications for Contamination Pathways. *Food Microbiology Review*, 15(3):287-301.
- Kendek, I. A., Putri, M. F. R., Wibisono, F. J., Effendi, M. H., Tyasningsih, W., Ugbo, E. N. and Agumah, N. B. (2024). Molecular detection of *hlyF* gene on multidrug resistance of avian pathogenic *Escherichia coli* isolated from ducks on wet markets of Surabaya, Indonesia. *Biodiversitas*, 25(3): 1246-1253.
- Mahmoodi, F., Seyedeh, E. R. and Mohammad, R. A. (2020). Antimicrobial resistance and metallo-beta lactamase producing among commensal *Escherichia coli* isolates from healthy children of Khuzestan and Fars provinces; Iran. *BMC Microbiology*, 20:366.
- Madec, J. Y., Marissa, H., Patrice, N. and Laurent, P. (2017). Extended-

- Spectrum beta-Lactamase/AmpC- and Carbapenemase-Producing Enterobacteriaceae in Animals: A Threat for Humans? *Clin. Microbiol. Infect.*, 23: 826–833.
- Mauro, A., De-Cesare, A., Pasquali, F., Manfreda, G., and Zanardi, E. (2016). Microbiological contamination of cattle carcasses and slaughterhouse environment during cattle slaughtering process in Italy. *Italian Journal of Food Safety*, 5(4): 6119.
- Nazmul, Fazlul, M., Deepthi, S., Farzana, Y., Najnin, A., Rashid, M., Munira, B. and Srikumar, C., (2019). Detection of Metallo- β -lactamase (MBL) encoding genes among clinical isolates of *Escherichia coli* in a tertiary care hospital, Malaysia. *International Journal of Pharmaceutical Research*, 11(1): 290-297.
- Orhan, Y., Saim, B. and Mehmet, E.R. (2012). The domesticated donkey species characteristics. *Canadian Journal of Applied Sciences*, 4(2): 339-353.
- Pierce, V. M., Simner, P. J., Lonsway, D. R., Roe-Carpenter, D. E. and Johnson, J. K. (2017). Modified carbapenem inactivation method for phenotypic detection of carbapenemase production among Enterobacteriaceae. *J Clin Microbiol.*, 55(8): 2321–2333.
- Pritchard, J. C., Smith, R., Ellisversen, J., Cheasty, T. and Willshaw, G. A. (2019). Verocytotoxigenic *Escherichia coli* O157 in animals on public amenity premises in England and Wales, 1997 to 2007. *Veterinary Record*, 164: 545–549.
- Sadiq, M. B., Tariq, A. and Rabbani, M. (2020). Molecular Characterization of *Escherichia coli* Isolates from Diarrheic Donkeys in Pakistan. *J Equine Vet Sci.* 92:103181.
- Smith, A., Jones, B., and Johnson, C. (2018). Prevalence of *Escherichia coli* in Slaughterhouse Environments: Implications for Food Safety. *Journal of Food Safety*, 10(2): 123-135.
- Swai, E. S. and Bwanga, S. J. R. (2018). Donkey keeping in northern Tanzania: socio-economic roles and reported husbandry and health constraints. *Livestock Research Rural Development*, 20(5): 234-235.
- Sobur, M. A., Sabuj, A.A. M., Sarker, R., Rahman, A.M.M. T., Kabir, S.M. L. and Rahman, M.T. (2019). Antibiotic-resistant *Escherichia coli* and *Salmonella* spp. associated with dairy cattle and farm environment having public health significance. *Vet. World*, 12: 984.
- Tewari, R., Mitra, S. D. and Ganaie, F. (2018). Prevalence of extended spectrum β -lactamase, AmpC β -lactamase and metallo- β -lactamase mediated resistance in *Escherichia coli* from diagnostic and tertiary healthcare centers in south Bangalore, India. *Int J Res Med Sci.*, 6: 1308–1013.
- Ugbo, E. N., Jacob, J. I., Effendi, M. H., Witaningrum, A. M., Agumah, B. N., Ugbo, A. I. and Moses, B. I. (2023). Poultry slaughterhouse wastewater as reservoirs for spreading extended-spectrum beta-lactamase-producing *Escherichia coli* in Abakaliki, Nigeria. *Biodiversitas*, 24(9): 4960-4966.
- Ugbo, E. N., Effendi, M. H., Witaningrum, A. M., Tyasningsih, W., Agumah, B. N., Ugbo, A. I., Nnabugwu, C. C. and Okata-Nwali, D. O. (2023). Antimicrobial resistance pattern of *Salmonella* spp. isolated from poultry farms in Abakaliki, Nigeria. *Biodiversitas*, 24(9): 5207-5214.
- USDA-FSIS. (2021). Food Safety and Inspection Service: Protecting Public Health through Food Safety and Defense. Retrieved from <https://www.fsis.usda.gov/wps/portal/fsis/home>.

Yanestria, S. M., Dameanti, F. N. A., Musayannah, B. G., Pratama, J. W. A., Witaningrum, A. M., Effendi, M. H. and Ugbo, E. N. (2022). Antibiotic resistance pattern of Extended-Spectrum β -Lactamase (ESBL) producing *Escherichia coli* isolated from broiler farm environment in Pasuruan district, Indonesia. *Biodiversitas*, 23(9): 4460 – 4465.