

## SHORT COMMUNICATION

Mycological Quality of Packaged Ginger Spiced Peanut Cake (*Kuli kuli*) Sold in Umuahia Metropolis, Abia State, Nigeria

Osaro-Matthew, R. C.\* and Itaman, V. O.

Department of Microbiology, College of Natural Science, Michael Okpara University of Agriculture, Umudike.

\*Correspondence author: [mrc.osaro@mouau.edu.ng](mailto:mrc.osaro@mouau.edu.ng)

**Abstract:** *Kulikuli* (peanut cake) is a known street snacks prone to fungal infestation due to poor production, handling and storage measures. The study aimed at evaluating the fungal quality of packaged and ginger spiced *kulikuli* consumed in Umuahia, metropolis, Abia State. Twenty four samples of *kulikuli* were sourced randomly from markets and supermarkets in Umuahia. Standard methods were adopted to assay the incidence of fungi, fungal load and moisture content of the *kulikuli* samples. Fungal isolates were identified based on their microscopic and cultural morphology. The moisture content ranged from  $4.4 \pm 0.03$  -  $6.3 \pm 0.2\%$ , and the total fungal load ranged from  $1.7 \pm 0.5$  -  $5.2 \pm 0.7 \times 10^5$  CFU/g. Forty eight morphologically distinct fungi that belong to 5 genera (*Mucor*, *Aspergillus*, *Penicillium*, *Rhizopus* and *Fusarium*) were isolated. *Aspergillus* had the highest percentage of occurrence (35.4 %) and the least occurring genus was *Fusarium* (8.3%). The least numbers of fungi 4 (8.3%) was recovered from packaged ginger spiced samples. This study suggest that *kulikuli* should be properly handled, packaged and spiced with appropriate quantity of herbs so as to reduce mould contamination of the snacks.

**Keyword:** *Kulikuli*; Fungi; Packaged; Ginger, Contamination.

## INTRODUCTION

Peanut cake commonly known as *kulikuli* is a product of groundnut. *Kulikuli* is an indigenous snacks widely consumed by old and young in the West African coast (Ezekiel *et al.*, 2011; Mupunga *et al.*, 2017). The process of production of *kulikuli* involves frying of the peanut (groundnut), milling into paste, extraction of oil and then frying or baking the chunk (Adjou *et al.*, 2012).

Just as the raw material (groundnut), *kulikuli* contains high protein and crude fat (Ezekiel *et al.*, 2011). It is an affordable street food which helps in meeting the essential nutritional needs of the teeming population of the West African coast (Mupunga *et al.*, 2017) especially Nigeria. *Kulikuli* is not just consumed by humans only, but also used in the formulation of poultry feed (Ademola *et al.*, 2015).

The high nutritive content and moderate moisture content of *kulikuli* made it a good substrate for the growth of fungi. Fungal contamination of *kulikuli* might take place during production, storage and sells. Among the fungal contaminants previously reported

on groundnut and groundnut products are fungi of genera *Aspergillus*, *Fusarium* and *Penicillium*. Most members of these genera are known producers of mycotoxins. Mycotoxins are toxins of fungal origin produced during secondary metabolism that are harmful to humans and animals when ingested (Ráduly *et al.*, 2020).

Due to market competition, some producers have devised a way of enhancing their end product acceptability by spicing their *kulikuli* with some herbs such as garlic, ginger, cinnamon etc. Some researchers have reported the ability of some of these herbs and spice to repress the growth of fungi (Agbebi *et al.*, 2013; Ayoade & Adegbite, 2016; Chuku, 2014). Some producers and vendors also properly package the *kulikuli* snacks in a sachet or container before sells. Despite these practices being common these days, there is paucity of data on its effect on the microbial quality of *Kulikuli*. Therefore this study is aimed at investigating the mycological quality of packaged and ginger spiced *kulikuli*.

## MATERIALS AND METHODS

### Sample collection:

A total of 24 samples of *kulikuli* (peanut cake) which comprised of 12 packaged (ginger spiced, 6 and non- ginger spiced, 6) and 12 unpackaged (ginger spiced, 6 and non- ginger spiced, 6), were sourced randomly from Umuahia metropolis markets and supermarkets.

### Enumeration, isolation and identification of fungi from *kulikuli* samples

*Kulikuli* samples were crushed and 10 g of each were homogenized in sterile distilled water (90 ml), and serially diluted ( $10^{-1}$  to  $10^{-5}$ ). One milliliter of the diluted aliquot was inoculated onto a molten Potato dextrose agar (PDA) supplemented with 0.01% streptomycin sulphate in duplicates using pour plate method. Incubation was carried out for 7 days at  $28 \pm 2^\circ\text{C}$ , developed distinct colonies were counted and the number of colonies from a plate with 30 – 300 colonies was multiplied by the inverse of the dilution factor to get the CFU/g. Fungal colonies morphologically different were subcultured thereafter identified macroscopically and microscopically using lactophenol cotton blue stain (Singh *et al.*, 1991; Samson *et al.*, 1995).

### Moisture content determination

The AOAC (2005) method was adopted in the determination of the moisture content. Five grams of crushed *kulikuli* samples were weighed and placed in crucibles of known weight. In a repeated 2 hourly, the crucibles containing the samples were placed in an oven preheated at  $105^\circ\text{C}$  and cooled in a desiccator until a constant weight was obtained. The moisture content was

calculated based on the reduction in weight and expressed as a percentage.

### Analysis of data

Analysis of data was carried out using SPSS 21.0. Calculated means of data were tested for significance by the Duncan multiple range test ( $P = 0.05$ ).

## RESULTS

The result obtained from this study showed that all samples were contaminated, the mean fungal count as presented in Table 1 showed that the un-packaged and un-spiced *kulikuli* samples had high fungal count ( $5.2 \pm 0.7 \times 10^5$  CFU/g) which was significantly ( $P < 0.05$ ) higher than the count obtained in packaged and ginger spiced *kulikuli* ( $1.7 \pm 0.5 \times 10^5$  CFU/g).

The morphological studies on the fungal isolates revealed that the isolates (48) were members of five genera; *Mucor*, *Aspergillus*, *Penicillium*, *Rhizopus* and *Fusarium*.

The percentage distribution of fungi revealed that *Aspergillus* was found in all samples and had the highest distribution of 35.4%, the least occurring genus was *Fusarium* with a distribution of 8.3%. *Mucor* and *Rhizopus* was not found in the packaged samples Table 2.

The moisture content of the *kulikuli* samples showed an elevated moisture content in the unpackaged samples, which the highest moisture content of  $6.3 \pm 0.20\%$  was obtained in the non- ginger spiced unpackaged sample, while the least moisture content of  $4.4 \pm 0.03\%$  was obtained in the ginger spiced packaged samples (Table 3). There was significant difference in the moisture contents of the samples ( $P < 0.05$ ).

**Table 1: Mean fungal count of *kulikuli* samples**

| Samples | Fungal count (x $10^5$ CFU/g) |
|---------|-------------------------------|
| PG      | $1.7 \pm 0.5^a$               |
| PUG     | $2.8 \pm 1.0^b$               |
| UPG     | $3.7 \pm 0.4^c$               |
| UPNG    | $5.2 \pm 0.7^d$               |

PG= Packaged ginger spiced *kulikuli*. PNG= Packaged non- ginger spiced *kulikuli*. UPG = Unpackaged ginger spiced *kulikuli*. UPNG = Unpackaged non- ginger spiced *kulikuli*.

Data with the same superscript alphabets are not significantly different at  $p \leq 0.05$  using the Duncan multiple range test.

**Table 2: percentage of fungal occurrence in *kulikuli* samples**

|                        | Samples<br>No. positive |         |          |          | Distribution No. (%) |
|------------------------|-------------------------|---------|----------|----------|----------------------|
|                        | PG                      | PNG     | UPG      | UPNG     |                      |
| <i>Mucor</i> spp       | -                       | -       | 3        | 5        | 8(16.7)              |
| <i>Aspergillus</i> spp | 3                       | 4       | 5        | 5        | 17(35.4)             |
| <i>Penicillium</i> spp | 1                       | 3       | 5        | 5        | 14(29.2)             |
| <i>Rhizopus</i> spp    | -                       | -       | 2        | 3        | 5(10.4)              |
| <i>Fusarium</i> spp    | -                       | 1       | 1        | 2        | 4(8.3)               |
| <b>Total No. (%)</b>   | 4(8.3)                  | 8(16.7) | 16(33.3) | 20(41.7) | 48(100)              |

PG= Packaged ginger spiced *kulikuli*. PNG= Packaged non-ginger spiced *kulikuli*. UPG = Unpackaged ginger spiced *kulikuli*. UPNG = Unpackaged non-ginger spiced *kulikuli*.

- = not detected

**Table 3: Moisture content of the *Kulikuli* samples**

| Samples | Moisture content (%)    |
|---------|-------------------------|
| PG      | 4.4 ± 0.03 <sup>a</sup> |
| PNG     | 4.8 ± 0.05 <sup>b</sup> |
| UPG     | 5.2 ± 0.10 <sup>c</sup> |
| UPNG    | 6.3 ± 0.20 <sup>d</sup> |

- PG= Packaged ginger spiced *kulikuli*. PNG= Packaged non-ginger spiced *kulikuli*. UPG = Unpackaged ginger spiced *kulikuli*. UPNG = Unpackaged non-ginger spiced *kulikuli*.

- Data with the same superscript alphabets are not significantly different at  $p \leq 0.05$  using the Duncan multiple range test.

## DISCUSSION

Mycological analysis of a street vended food gives us insight into the number and types of fungi present therein, which depicts the food quality and possible risk associated with its consumption. All the *kulikuli* samples analyzed were contaminated with fungi which is in agreement with the reports of Adetunji *et al.* (2018), Ayoade & Adegbite (2016) and Ezekiel *et al.* (2011) who recorded the presence of fungi in all the groundnut and groundnut product samples studied.

The mean fungal count was highest in the unpackaged non-ginger spiced samples, this might be attributed to the fact that these *kulikuli* samples were not spiced with ginger which have been associated with antifungal properties (Agbebi *et al.*, 2013, Chuku, 2014). Also the fact that these samples were not packaged might have exposed them to fungal spores and conidia. The least fungal count was obtained from the packaged

ginger spiced *kulikuli* samples, this means that the ginger repressed the growth of fungi and the pack protected the samples from contamination. Generally, the fungal counts obtained from all the *kulikuli* samples were above the  $10^3 - 10^4$  CFU/g acceptable limits in dried food (ICMSF, 2018), but none of these samples met with the criteria, this heavy contamination might have occurred at the pre-processing stage or the processing stage (Ayoade & Adegbite, 2016).

The fungal strains obtained from this study belong to the genera *Mucor*, *Aspergillus*, *Rhizopus*, *Penicillium* and *Fusarium*. These genera have been reported among the commonly associated microflora of groundnut and its products (Ezekiel *et al.*, 2011; Odeniyi *et al.*, 2019). Most species of *Aspergillus*, *Penicillium* and *Fusarium* are mycotoxigenic, mycotoxins when ingested are nephrotoxic, hepatotoxic, cytotoxic, genotoxic, carcinogenic and teratogenic (Ráduly *et al.*, 2020).

*Mucor* and *Rhizopus* obtained from this study are known for their biodeterioration ability (Mupunga *et al.*, 2017) which will adversely affect the nutritive quality and the organoleptic properties of the *kulikuli*. Also some strains of *Rhizopus* are toxigenic (Othman *et al.*, 2018) hence making the presence of these genera in food a thing of concern.

The percentage occurrence of the various fungi showed that *Aspergillus* was found in all samples and had the highest percentage distribution of 35.4% followed by *Penicillium*. *Aspergillus* is notorious for its high adaptability and dispersibility, this might explain its dominance amongst the samples. In concordance with this report is the result of Adetunji *et al.* (2018) where *Aspergillus* was the dominant fungi and had a percentage distribution of 60% in groundnut samples. *Mucor* and *Rhizopus* were isolated only from the unpacked samples, this is an indication that the contamination must have occurred due to improper handling during storage or sales. *Fusarium* was the least occurring fungi genera from this study, this is in agreement with the report of Adeoye *et al.* (2020) who obtained 5% distribution of genus *Fusarium* in Kulikuli. And contrary to that of Odeniyi *et al.* (2019) where *Fusarium* occurred at 33% in Kulikuli, this disagreement might be as a result of variation in production procedure and environmental factors.

The moisture content obtained in this study is consistent with the report of Odeniyi *et al.* (2019) who obtained a moisture content of

4.55 – 6.3 % range in *kulikuli* samples from two south west (Lagos and Ogun State) Nigerian geopolitical zone. The moisture content in the unpackaged *kulikuli* snacks exceeded the acceptable moisture content limit of > 5% for nuts and nuts products storage (Adetunji *et al.*, 2018), this might have resulted from the unpackaged samples absorbing moisture from its surrounding environment. According to Magan *et al.* (2010) moisture content is critical in the accumulation of fungi and mycotoxin in food.

## CONCLUSION

The *kulikuli* samples studied were all contaminated with fungi, this call for concern as proliferation of these fungi in a favorable condition might make the snacks a medium for food intoxication. The study revealed packaged and ginger spiced *kulikuli* samples had reduced load of fungi ( $1.7 \pm 0.5 \times 10^5$  CFU/g) compared to the unpackaged and un-spiced *kulikuli* samples ( $5.2 \pm 0.7 \times 10^5$  CFU/g). Fungal counts obtained from all samples were above the International Commission for Microbiological Specification for Food (ICMSF) acceptable limits of  $10^3$  -  $10^4$  CFU/g of groundnut products sample. Several moulds were isolated with the most occurring genus being *Aspergillus*, many of these moulds have been implicated in the production of mycotoxins. These findings implies that consumers are possibly exposed to mycotoxins, hence a need for public health interventions.

## REFERENCES

- Ademola, S. G., Shittu, M. D. Osanyande, M. O. & Omididi, O. E. (2015). Effects of mycotoxin detoxifier supplementation to contaminate Groundnut cake on the growth performance organs and serum enzymes of two broiler strains. *International Journal of Agriculture Innovations and Research*, 4(2): 2872.
- Adetunji, M. C., Alika, O. P., Awa, N. P., Atanda, O. O. & Mwanza, M. (2018). Microbiological quality and risk assessment for aflatoxin in groundnut and roasted cashew nuts meant for human consumption. *Journal of Toxicology*, 1308748, 1-11.
- Adeyo, A. O., Adeyinka, O.O., Ajala, A.S. & Ojo, M. A. (2020). Microbiological assessment of selected groundnut based snacks

- kulikuli and donkwa in Ogbomoso and Ilorin, Nigeria. *Acta Scientific Nutritional Health*, 4(4), 67-71.
- Adjou, E. S., Yehouenou, B. & Spssou, C. M. (2012). Occurrence of mycotoxins and associated mycoflora in peanut cake product (kulikuli) marketed in Benin. *African Journal of Biotechnology*, 11(76), 14354.
- Agbebi, O. T., Ilesanmi, A. I. & Abdurahem, I. (2013). Additive potentials of *Zingiber officinale* on the aflatoxin secreted in waste mould feed fed to *Clarias gariepinus* (Burchell 1822). *International Journal of Biological and Chemical Sciences*, 7(1): 236 – 346.
- AOAC (Association of Official Analytical Chemist) (2005). *Official methods of analysis*. 18<sup>th</sup> Ed. Washunton DC pp. 949.
- Ayoade, F. & Adegbite, T. D. (2016). Microbiological screening of street vended groundnut kulikuli and natural spices for reducing microbial contamination in the food snack. *International Journal of Chemical Sciences*, 10(6): 2677- 2691.
- Chuku, E. C. (2014). Antifungal properties of ginger (*Zingiber officinnale*) and garlic (*Allium sativum*) on smoked cat fish mycroflora. *Nigerian Journal of Mycology*. 6: (48 -57).
- Ezekiel, C. N., Anokwuru, C. P., Fari, A., Olorunfemi, M. F., Fadairo, O., Ekeh, H. A., Ajoku, K. Gbuzue, N. & Akinsanmi, F. (2011). Microbiological quality and proximate composition of peanut cake (kulikuli) in Nigeria market. *Academia Arena*, 3(4), 103 – 111.
- ICMSF (International Commission for Microbiological Specification for Food) (2018). *Microorganisms of Food* 7. Microbiological testing in food safety management. 2<sup>nd</sup> Ed. Springer.
- Magan, N., Aldred, D., Mylon, K. & Lambert, R.J.W. (2010). Limiting mycotoxins in stored food. *Food Additives and Contaminants Part A*. 27, 644 – 650.
- Mupunga, I., Mngqawa, P. & Katerere, D. R. (2017). Peanuts, aflatoxin and under nutrition in children in Sub-saharan Africa. *Nutrients*. 9, 1287 - 1299.
- Odenyi, O. O., Ojo, C., Adebayo-Tayo, B. & Olasenhinde, K. (2019). Mycological toxicogenic and nutritional characteristics of some vended groundnut and groundnut products from three Northern Nigeria ecological zones. *African Journal of Biomedical Research*. 22, 65 -71.
- Othman, G. A., Ahmed, D. A., Alghamdi, H. A. & Radwan, A. M. (2018). Antifungal activity of *Syzygium aromaticum* (Dianthus) against toxigenic *Rhizopus stolonifer* and its immunomodulatory effects in aflatoxin-fed mice. *Tropical Journal of Pharmaceutical Research*, 17,10.
- Ráduly, Z., Szabó, L., Madar, A., Pósci, I. & Csernoch, L. (2020). Toxicology and medical agents of *Aspergillus*-derived mycotoxins entering the feed and food chain. *Frontier in Microbiology*, 10: 2908.
- Samson, R. A., Hoekstra. E. S., Frisvad, J. S. & Filtenborg, O. (1995). Methods to detection and isolation of food-borne fungi, pp 235 -242. In Samson, R. A., Hoekstra. E. S., Frisvad, J. S., Filtenborg, O. *Introduction to food borne fungi*. 4<sup>th</sup> Ed. Central Bureau Voor Scjimmel Cultures. The Netherland pp. 235-242.
- Singh, K., Frisvad, J. C., Thrane, U. & Mathu, S. B. (1991). *An illustrated manual of identification of some selected food borne Aspergilli, Fusaria, Penicillia, and their mycotoxins*. Heller u Denmark, Danish government. Institute of Seed Pathology for Developing Countries. Pp 132.