

N. J. Microbiology, 2(2), 1982, pp 141 – 146
© The Nigerian Society for Microbiology 1982.

THE VAGINAL BACTERIA FLORA IN PREGNANT AND NON-PREGNANT WOMEN

By

BEATRICE NGOZI ANYANWU

Lecturer, Department of Microbiology,
University of Nigeria, Nsukka, Nigeria.

FELICIA OBIOMA CHIEDOZIE

Department of Microbiology,
University of Nigeria, Nsukka, Nigeria.

ABSTRACT

The bacterial flora of the vagina was assessed in healthy Nigerian women (twenty pregnant and twenty-four non-pregnant women). They all harboured many types of microorganisms in their vagina. The mean number of microbial types isolated was five for the pregnant and three for the non-pregnant women. *Staphylococcus* was detected in 18 of the pregnant women and 16 of the non-pregnant women. Amongst the anaerobes *Peptostreptococcus* was isolated in 14 of the pregnant women and 6 of the non-pregnant women. Lactobacilli in 7 and 12 respectively. Bacteroides was not isolated from the vagina of pregnant women but from 50% of the vaginal swabs obtained from the non-pregnant women.

Statistically, the difference between the bacterial flora of the pregnant and non-pregnant women investigated is not significant.

INTRODUCTION

Several laboratories have studied the microbial flora of the female genital tract. Gorbach, Menda, Thadepalli and Keith (1973) found the most common anaerobes to be bacteroides followed by peptostreptococci. Corbishley (1977) found that both vagina and cervix harboured many types of microorganisms in common but that the concentrations of the organisms differed between the sites. The purpose of this report is to compare the bacterial flora of pregnant women with that of non-pregnant women.

MATERIALS AND METHODS

Cultures were obtained from the vagina of 20 non-pregnant women attending the Gynecology Out-Patient Clinic at the University of Nigeria Teaching Hospital, Enugu; 20 pregnant women attending prenatal clinic at the University of Nigeria Teaching Hospital Enugu (Department of Gynecology and Obstetrics) and four non-pregnant students of the University of Nigeria Nsukka. Five of the pregnant women were in their first trimester of pregnancy and the rest were in their mid or last trimester of pregnancy. The women were all healthy and were between 18 and 40 years old and the average age was 28. Women receiving birth control pills or antibiotics and women with signs of genital infections were excluded from this study.

SAMPLING TECHNIQUE

Vaginal secretions were obtained during pelvic examination. Cotton-tipped applicators were inserted and then rotated in the posterior fornix. Care was taken to avoid any contact of the swab with the external genitals, or other sources of contamination. Similar swabs were also obtained from the four students with the help of a medical doctor at the University of Nigeria Medical Centre.

The swabs were dipped into Stuart Transport Medium (Oxoid) and delivered to the Microbiology Laboratory of the University of Nigeria Nsukka, within four hours of collection, for the processing. Each of the swabs was inoculated onto five primary culture media. The media used were:

(1) blood agar (oxid) (2) MacConkey agar (oxid) (3) blood agar with neomycin (4) tomato juice agar (oxid) and (5) chocolate agar. Plates (1) and (2) were incubated aerobically, plates (3) and (4) anaerobically and plate (5) was incubated in 10% carbon dioxide for the isolation of *Neisseria gonorrhoea*.

The anaerobic system was obtained by using Fildes and McCintosh jars from Messrs Baird and Tatlock. The jars were evacuated and replaced with oxygen-free CO₂ (10%) and hydrogen (90%). The anaerobic jars were incubated for 72 hours at 37°C. To check the system, viable *Pseudomonas aeruginosa*, an obligate aerobe was included. This will fail to grow if true anaerobiosis was established. An environment of 10% CO₂ was achieved by burning a candle stick in a closed container in which plate (5) was incubated. Aerobic Culture were examined after 24 hours and again at 48 hours incubation.

All colonial types on each medium were investigated further by microscopy, biochemical and physiological tests, and divided into genera or species by routine laboratory methods, (Cruickshank, Duguid, Marmion, and Swain 1975), to give the categories listed in Table III.

RESULT

Most specimens yielded polymicrobial growth. Total isolates were 99 from 20 pregnant women and 77 from 24 non-pregnant women. The mean number of isolates per specimen being five for the pregnant women and three for the non-pregnant women. (See Table I).

Seventy five percent of the pregnant women showed positive cultures for both aerobes and anaerobes. Twenty five percent showed positive cultures for aerobes only while none showed positive cultures for anaerobes only. For the non-pregnant women 58.3% showed positive cultures for both aerobes and anaerobes, 41.7% showed positive cultures for aerobes only and none showed positive cultures for anaerobes only. (See Table II).

Statistical analysis of the data showed no significant difference in the vaginal flora of pregnant and non-pregnant women at 5% error.

(No. of Microbial
Isolates per
specimen.

0
1
2
3
4
5
6
7
8

Total 99

No. of positive
cultures

Aerobes &
anaerobes

Aerobes only

Anaerobes only

Total

TABLE I

FREQUENCY OF MULTIPLE ISOLATES OF MICROBIAL SPECIES FROM THE VAGINA OF PREGNANT AND NON-PREGNANT WOMEN.

PREGNANT		NON-PREGNANT	
No. of Microbial Isolates per specimen.	No. of specimens.	No. of Microbial Isolates per specimen.	No. of specimens.
0	0	0	0
1	0	1	2
2	1	2	5
3	3	3	10
4	6	4	3
5	5	5	2
6	4	6	1
7	1	7	1
8	1	—	—
Total 99	20	77	24

TABLE II

INCIDENCE OF POSITIVE CULTURES FROM THE VAGINA OF PREGNANT AND NON-PREGNANT WOMEN.

PREGNANT			NON-PREGNANT	
No. of positive cultures.		Percent of total	No. of Positive cultures.	Percentages of total
Aerobes & anaerobes	15	75%	14	58.3
Aerobes only	5	25%	10	41.7
Anaerobes only	0	0%	0	0%
Total	20	100%	24	100%

TABLE III
FREQUENCY OF ISOLATES

Organisms	Pregnant Women	Non-Pregnant Women	Total	Percentage Occurrence
	(20)	(24)	(44)	
<i>Aerobes:</i>				
<i>Staphylococcus</i> sp.	18	16	34	77.4
<i>Strep. faecalis</i>	4	6	10	25.0
<i>Micrococcus</i> sp.	4	2	6	17.7
<i>Diphtheroids</i>	12	2	14	31.9
<i>E. coli</i>	8	6	14	31.9
<i>Klebsiella pneumoniae</i>	6	2	8	18.2
<i>Proteus</i> sp	2	0	2	4.6
<i>Lactobacillus</i> sp	4	8	12	27.3
<i>Bacillus</i> sp	4	4	8	18.2
<i>Anaerobes:</i>				
<i>Peptococcus</i> sp	8	12	20	45.5
<i>Peptostreptococcus</i> sp.	14	6	20	45.5
<i>Clostridium</i> sp	2	0	2	4.6
<i>Lactobacillus</i> sp	3	4	7	15.9%
<i>Bacteroides</i> sp	0	12	12	27.3%

The aerobes isolated were *Staphylococcus* sp (*S. epidermidis* and *S. Aureus*). *Streptococcus* (*S. faecalis* only) *Micrococcus* Sp. *Diphtheroids*, *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus* and *Bacillus*. The anaerobic isolates were *Peptococcus* *Peptostreptococcus*, *Bacteroides*, *Clostridium* and *Lactobacillus*. The genus *Lactobacillus* covers both anaerobic and facultative isolates. *Neisseria gonorrhoea* was not isolated in any of the samples.

Table III shows the frequencies of the isolates. (See Table III).

DISCUSSION

The study shows no statistical significant difference between pregnant and non-pregnant vaginal flora. This could be due to the smallness of the sample size. But there is a difference both in types and relative concentrations of the bacteria isolated. The vaginal flora may vary depending on factors such as age, hormonal influence, pregnancy, use of contraceptives, use of antibiotics and state of health of the individual (Hurley, *et al* (1974). Bartlett, *et al* (1977)

have pointed out constant fluctuations in types and relative concentrations of the bacteria isolated. The results of this study are in line with those of deLouvois (1977) who found that the vaginal flora of non-pregnant women is dominated by *Staphylococcus aureus*, *Corynebacterium* and *Lactobacillus*. The results of this study are also in line with those of deLouvois (1977) who found that the vaginal flora of pregnant women is dominated by *Staphylococcus aureus*, *Corynebacterium* and *Lactobacillus*.

The results of this study are also in line with those of deLouvois (1977) who found that the vaginal flora of pregnant women is dominated by *Staphylococcus aureus*, *Corynebacterium* and *Lactobacillus*. The results of this study are also in line with those of deLouvois (1977) who found that the vaginal flora of pregnant women is dominated by *Staphylococcus aureus*, *Corynebacterium* and *Lactobacillus*.

The absence of *Neisseria gonorrhoea* amongst the isolates is not surprising in view of the range of socio-economic factors involved.

Of the anaerobes, *Peptococcus* and *Peptostreptococcus* were the most common, accounting for 50% for the non-pregnant women and 45.5% for the pregnant women. *Streptococcus faecalis* was isolated 19 times in 20 samples.

The isolation of *Lactobacillus* is noteworthy. In this study, it was isolated 15 times in 44 samples (34%). Corbier *et al* (1973), isolated 15 *Bacteroides* in 100 samples (15%). deLouvois *et al* (1977) isolated 15 *Clostridium* in 100 samples (15%). Swenson *et al* (1977) isolated 15 *Clostridium* in 100 samples (15%).

The difference in the incidence of the bacteria isolated is significant. 60% of the bacteria isolated cover organisms found in both pregnant and non-pregnant women.

have pointed out that the vagina harbours an unstable bacterial population that is subjected to constant fluctuations based on well defined hormonal influences and ill-defined extraneous factors. The results of this study have confirmed these views. Here, there are fluctuations both in types and relative concentrations of the bacteria isolated from non-pregnant and pregnant women.

Staphylococcus possessed the highest percent incidence 90% and 66.6% in pregnant and non-pregnant females respectively, 73% of these Staphylococci were *S. epidermidis* and 27% *S. aureus*. Corbishley (1977) observed 89% and 73% coagulase negative staphylococci in vaginal and cervical swabs respectively. 17% coagulase positive staphylococci were observed in each site. deLouvois, *et al* (1975) observed 66.1% and 4.6% *S. epidermidis* and *S. aureus* respectively in pregnant women.

Lactobacilli have been thought to be preponderant in the vagina. In this investigation they were found in 35% and 50% of the vaginal swabs of pregnant and non-pregnant women respectively. Hurley *et al* (1974) pointed out that in children and women of menopausal age the predominating bacteria are the cocci, while lactobacilli predominated during pregnancy and at puberty. The incidence in both groups was 43.2%. Other surveys show incidences of lactobacilli varying between 0% and 82%: Morris and Morris (1967) 49.1%; Gorbach *et al* (1973) 73%; deLouvois, *et al* (1975) 81.8%; Ohm and Galask, (1975) 75%; Swenson, *et al.* (1973) 0.0%. Swenson *et al.* (1973) studied the anaerobic bacterial infection of the female genital tract, and recovered all the anaerobes found in the study except *Lactobacillus*. Saigh, *et al.* (1977) observed the inhibitory effect of endocervical flora mainly lactobacilli on the growth of *N. gonorrhoea*. They found that when the endocervical flora is low or absent the pathogen increased in number.

The absence of *N. gonorrhoea* in this study is explained by the fact that this study was confined to mainly married middle class women. In a separate study the incidence of *N. gonorrhoea* amongst Nigerian women will be reported. The study will cover more samples and a wider range of socio-economic groups in Nigeria.

Of the anaerobes isolated *Peptostreptococcus* occurred more in pregnant women than in non-pregnant women, 70% and 25% respectively. *Peptococcus* was 40% for the pregnant and 50% for the non-pregnant. In other studies, (Gorbach *et al* 1973) isolated 33% and 10% *Peptostreptococcus* and *Peptococcus* respectively. Swenson *et al* (1973) isolated *Peptostreptococcus* sp. 19 times and *Peptococcus* sp. 21 times in 91 infections of the genital tract.

The isolation of bacteroides 50% in non-pregnant women and none in pregnant women is noteworthy. In other studies on genital tracts, bacteroides were isolated. Swenson *et al.* (1973), 66.6%; Corbishley (1977), 24% from vaginal swabs and 20% from cervical swabs. Gorbach *et al* (1973), 57%. The women in these studies were not pregnant. DeLouvois *et al* (1975) isolated 15 bacteroides spp. from 280 unselected pregnant women (5.4%), the most reasonable explanation of the absence of bacteroides is probably the small number sampled, but the incidence of bacteroides is definitely less in pregnant women. *Proteus* and *Clostridium* were isolated (10% each) in pregnant women but none was isolated in non-pregnant females. In other studies both genera have been isolated both in healthy and in infected female genital tracts. Gorbach *et al.* (1973), 10% and 17%; Corbishley (1977), 3% and 7% for vagina and 1% and 7% for cervix, deLouvois *et al.* (1975), 6.1% and 0.0% *Proteus* and *Clostridium* respectively. Swenson *et al.* (1973) studying the anaerobic bacterial infections of the female genital tract isolated *Clostridium* six times from 91 infections of the female genital tract. On the whole, the incidence of the two genera in vagina is low.

The difference in the incidence of diphtheroids in pregnant and non-pregnant women is significant. 60% for pregnant and 8.3% for non-pregnant women. Diphtheroid bacterial here, cover organisms suspected but not confirmed as *Corynebacterium*. Percentage occurrence in both pregnant and non-pregnant women was 31.9%. In other works, deLouvois *et al.* (1975)

observed 83.4% incidence of Corynebacteria in pregnant women. Corbishley (1977) observed 69% and 64% diphtheroids from vaginal and cervical swabs respectively.

Most of the anaerobes and some of the aerobes isolated are frequently involved in genital tract infections suggesting that the normal flora of vagina is potentially pathogenic.

LITERATURE CITED

- BARTLETT J.G. ONDERDON A.B. BRUDE E. GOLDSTEIN C. ANDERKA M., and McCORMACK, W.M. 1977. Quantitative bacteriology of the vaginal flora. *Journal of Infectious Diseases* 136: 271 – 277.
- CORBISHLEY C.M. 1977. Microbial flora of the vagina and cervix., *Journal of Clinical Pathology* 30: 745 – 748.
- GORBACH, S.L., MENDA K.B. THADEPALLI H. and KEITH, L. 1973. Anaerobic microflora of the cervix in healthy women. *American Journal of Obstetrics and Gynecology*. 177: 1053 – 1055.
- CRUICKSHANK R., DUGUID J.P. MARMION B.P. and SWAIN R.H.A. 1975. Identification of Specific Microbes. In *Medical Microbiology* 12th Edition Vol. 2 Churchill Livingstons, Edinbury.
- de LOUVOIS J., HURLEY R. and STANLEY V. C. 1975. Microbial flora of the Lower genital tract during pregnancy: relationship to morbidity. *Journal of Clinical Pathology* 28, 731 – 735.
- HURLEY R.S., STANLEY V. C., LEASK B.G.S. and deLOUVOIS J. 1974. Microflora of vagina during pregnancy. In *the Normal Microbial flora of Man*. pp. 155-182. Edited by Skinner F. K. and CARR J. G. Society of Applied Bacteriology, New York, Academic Press.
- MORRIS C. A. and MORRIS D.E. 1967. Normal Vaginal Microbiology of women of childbearing age in relation to the use of Oral Contraceptives and Vaginal tampons. *Journal of Clinical Pathology* 20. 636 – 640.
- OHM, M. J. and GALASK R.P. 1975. Bacterial flora of the Cervix from 100 pre hysterectomy patients. *American Journal of Obstetrics and Gynecology* 122: 683 – 687.
- SAIGH J. H., SANDERS C. C. and SANDERS E. J. 1977. Inhibition of *Neisseria gonorrhoeae* by Aerobic and Facultative anaerobic components of the Endocervical Flora. Evidence for the protective effect against Infection. *Immun.* 19: 704 – 710.
- SWENSON R. M., MICHAELSON T.C., DALY M. J., FACOG and SPAULDING E. H. 1973. Anaerobic bacterial infection of the female genital tract. *American Journal of Obstetrics and Gynecology* 42, 538 – 541.

EXPERIM

West
(Nairobi sh
were demon

Dugbe vir
Further vir
culicoides.
Nairobi she

Morph
jam (Dav
isolated in
African D
ty for this

Virus: Th
Dugbe m
passages i

Animals:
tive for s
thics, bat
imals wer
brain pas
rise in re
leucocyte
blood in
tested fo
Neutralis
method
as differ

Nor
Two she
1 and 2